

# Ultrasound training recommendations for medical and surgical specialties

Second edition

Board of the Faculty of Clinical Radiology  
The Royal College of Radiologists

# Contents

<b>Foreword</b>	<b>3</b>
<b>1 Introduction</b>	<b>4</b>
<b>2 Aims and principles</b>	<b>5</b>
<b>3 Training recommendations</b>	<b>8</b>
3.1 <i>Theoretical training</i>	8
3.2 <i>Practical training</i>	8
<b>4 Continuing professional development</b>	<b>10</b>
<b>References</b>	<b>11</b>
Appendix 1. Urological ultrasound	12
Appendix 2. Gynaecological ultrasound	18
Appendix 3. Gastrointestinal ultrasound	24
Appendix 4. Vascular ultrasound	29
Appendix 5. Breast ultrasound	33
Appendix 6. Thoracic ultrasound	37
Appendix 7. Cranial ultrasound in infants	40
Appendix 8. Focused emergency ultrasound	44
Appendix 9. Ultrasound training for critical/intensive care	48
Appendix 10. Musculoskeletal ultrasound	52
Appendix 11. Recommended theory syllabus	58

# Foreword

In 2005, when The Royal College of Radiologists (RCR) produced the initial document on *Ultrasound Training Recommendations for Medical and Surgical Specialties*, it recognised ultrasound as an evolving technology with wide application throughout medical and surgical practice.

Since 2005, there has been continuing growth in the availability of ultrasound and with it an acknowledgement that it can play an ever-increasing role in the diagnosis and management of patients.

It is, therefore, timely that the RCR updates this valuable guidance resource as part of its commitment to the development of high-quality imaging across the gamut of modalities. The original document, *Ultrasound Training Recommendations for Medical and Surgical Specialties*, has now been withdrawn.

The RCR wishes to ensure that access to high-quality ultrasound imaging continues to improve. For this to happen, ultrasound must be provided by properly trained and committed practitioners using appropriate quality ultrasound equipment.

In the UK, radiologists and sonographic practitioners have traditionally provided such a service from centralised departments. However, there is a growing need to provide ultrasound in other settings; for instance, in the community or by the patient's bedside in critical care.

By adopting the recommendations in this document, those involved in ultrasound training can ensure that wherever ultrasound is performed the patient receives the best possible service.

I would like to thank Rani Thind for her work in leading the revision of this advice and the following members of the working party for their contributions and advice:

- Dr Paul Dubbins, Clinical Radiologist
- Dr Fergus Gleeson, Clinical Radiologist
- Dr Alan Sprigg, Clinical Radiologist
- Dr Paul Sidhu, Clinical Radiologist
- Dr Richard McWilliams, Clinical Radiologist
- Dr Mike Weston, Clinical Radiologist
- Dr Marcus Harbord, Gastroenterologist

Thanks are also due to the following for their expert advice on the content: Dr Simon Jackson, Professor Andy Evans, Dr Charles Wakeley, Dr Eugene McNally, Dr Phil Cook, Dr Roger Bury and Dr Peter Wylie.

**Dr Pete Cavanagh**

Vice President and Dean, Faculty of Clinical Radiology  
The Royal College of Radiologists

# 1 Introduction

- 1.1 High-quality ultrasound services are provided by properly trained and committed practitioners using appropriate quality ultrasound equipment. In the UK, radiologists and sonographic practitioners have traditionally provided such a service from centralised departments of clinical radiology where equipment and manpower can be used cost-effectively.
- 1.2 Departments of clinical radiology may have difficulty responding to these demands primarily because of the national shortages of radiologists and sonographic practitioners. It is therefore essential that alternative methods of service delivery are considered. These may include the involvement of other professional groups in addition to greater investment in clinical radiology departments.
- 1.3 Medical specialists other than radiologists are increasingly wishing to undertake ultrasound examinations on patients referred to them for their clinical opinion as a direct extension of their clinical examination. This may take place in the outpatient department, on the wards, and in the assessment of emergency patients.

Clinicians are also using ultrasound to assist in practical procedures such as central line insertion. A separate document is aimed at providing guidance for focused ultrasound training.<sup>1</sup>

This should read some European and UK boards have incorporated ultrasound experience into clinical training and accreditation.

- 1.4 Radiologists have the skills, experience and commitment to provide guidelines for training of medical non-radiologists and hence influence the quality of service provided for the better. The RCR believes that this approach, of interdisciplinary co-operation, serves best the interests of patients.
- 1.5 Training of medical non-radiologists should be adequately funded and planned so that there is minimal adverse impact on the service provided to patients and the ability of clinical radiology departments to train clinical radiologists and sonographer practitioners.
- 1.6 This document makes recommendations for ultrasound training in the following areas:
  - Urological ultrasound
  - Gynaecological ultrasound
  - Gastrointestinal ultrasound
  - Vascular ultrasound
  - Breast ultrasound
  - Thoracic ultrasound
  - Cranial ultrasound in infants
  - Focused emergency ultrasound
  - Intensive care ultrasound
  - Musculoskeletal ultrasound.

## 2 Aims and principles

- 2.1 The medical use of ultrasound remains highly operator-dependent in spite of advances in technology, and the interests of the patient are best served by the provision of an ultrasound service which offers the maximum clinical benefit and optimal use of resources; that is, with appropriately trained personnel using equipment of appropriate quality.
- 2.2 All those who provide an ultrasound service are ethically and legally vulnerable if they have not been adequately trained. National Health Service (NHS) trusts and health boards in the UK, which provide professional indemnity to practitioners, are unlikely to be able to mount any defence to an action brought against an untrained practitioner. Similarly, the professional defence organisations are unlikely to be successful in mounting a defence against a claim for negligence should an error of diagnosis be made by an untrained practitioner of ultrasound. Advisory guidelines for training in ultrasound provided by the RCR will establish the principles to allow appropriate bodies to provide professional indemnity by setting out training and continuing professional development (CPD) requirements. Trusts, health boards, purchasing commissioners and patients should be aware of the requirements for training.
- 2.3 An appropriate level of training in ultrasound is one that allows for the provision of a safe and effective ultrasound service. This may be a purely diagnostic, predominantly interventional, or a clinically focused service. Departments of clinical radiology would normally provide all of these services, but it would be expected that other medical practitioners would deliver only those aspects of ultrasound particularly relevant to their clinical practice. Nonetheless, the training for medical non-radiologists should be to the same standard as those for radiologists, albeit restricted to the relevant and particular area of their clinical expertise. Whereas radiological training provides for the practice of ultrasound across a broad range of medical and surgical specialties, NHS trusts, health boards, purchasing commissioners and patients should be aware of the differences in the comparative depth and breadth of training, and hence ultrasound skills, between trained radiologists and trained medical non-radiologists.
- 2.4 The RCR has worked closely for many years with the Royal College of Obstetricians and Gynaecologists (RCOG) to ensure adequate training in obstetric ultrasound and obstetric ultrasound is not covered in this publication. It is also recognised that the RCOG has its own training module for ultrasound imaging in the management of gynaecological conditions and a number of radiologists act as preceptors for this. However, a syllabus for gynaecological ultrasound has been included as there are other groups (such as general practitioners) who might wish to train in this area.
- 2.5 The European Federation of Societies for Ultrasound in Medicine and Biology (EFSUMB) has proposed minimal training requirements for the practice of medical ultrasound in Europe.<sup>2</sup> These are supported by the RCR and the British Medical Ultrasound Society.
- 2.6 Three levels of minimum training requirements are proposed in this document.

### 2.6.1 *Level 1*

Practice at this level would usually require the following abilities:

- To perform common examinations safely and accurately
- To recognise and differentiate normal anatomy and pathology
- To diagnose common abnormalities within certain organ systems
- To recognise when a referral for a second opinion is indicated
- To understand the relationship between ultrasound imaging and other diagnostic imaging techniques.

Within most medical specialties, the training required for this level of practice would be gained during conventional postgraduate specialist training programmes. In the UK, this level of training would equate to the end of basic training in ultrasound of radiology specialist registrars (SpRs) in year 3/4 of training. It would also be equivalent to a holder of, for example, the RCOG special skills training module in gynaecological ultrasound imaging.

### 2.6.2 *Level 2*

Practice at this level would usually require most or all of the following abilities:

- To accept and manage referrals from Level 1 practitioners

- To recognise and correctly diagnose almost all conditions within the relevant organ system and to have sufficient understanding of ultrasound depiction of pathology to optimise the referral of the patient if the condition falls outside of the practitioner's skills
- To perform common non-complex ultrasound-guided invasive procedures
- To teach ultrasound to trainees and Level 1 practitioners
- To conduct some research in ultrasound

The training required for this level of practice would be gained during a period of subspecialty training which may either be within or after the completion of a specialist training programme. This would equate to the level of training in radiology at the time of acquiring the Certification of Completion of Training (CCT), assuming that part of the fifth year of subspecialty training had involved ultrasound.

### 2.6.3 Level 3

This is an advanced level of practice, which includes some or all of the following abilities:

- To accept tertiary referrals from Level 1 and Level 2 practitioners
- To perform specialised ultrasound examinations
- To perform advanced ultrasound-guided invasive procedures
- To conduct substantial research in ultrasound
- To teach ultrasound at all levels
- To be aware of and to pursue developments in ultrasound.

In the UK, this would equate to a consultant radiologist with a subspecialty practice which includes a significant commitment to ultrasound.

- 2.7 The boundaries between the three levels are difficult to define precisely and the above should only be regarded as a guide to different levels of competence and experience. In the detailed syllabuses attached to this document in Appendices 1–9 an attempt is made to indicate more specifically the type of experience required for each level of training. Training in musculoskeletal ultrasound does not lend itself easily to 'levels' of training and instead a 'modular' approach is recommended (Appendix 10).
- 2.8 The training of medical non-radiologists should foster relationships between radiological and non-radiological medical practitioners so that mutual support continues beyond the initial training period. Ideally a radiologist would continue to act as a mentor for a medical non-radiologist undertaking ultrasound after their training is completed. In addition, regular multidisciplinary meetings should continue to ensure an integrated approach to any further imaging that may be required.
- 2.9 A system for recording the results of any ultrasound examination in the patient's record is mandatory. The permanent recording of images, where appropriate, is also mandatory for the purposes of correlative imaging, future comparison and audit. The preferred option is through the hospital radiology information system (RIS)/picture archive and communications systems (PACS) equipment, enabling other clinicians to access the images and report.
- 2.10 Knowledge of the appropriate use and integration of other imaging techniques, as well as the clinical and economic impact of ultrasound on the demand for other imaging, should be acquired.
- 2.11 The requirement to deliver training for medical non-radiologists must acknowledge the time commitment of the trainer and trainee, the provision of funding, the content and practicability of the syllabus and the availability of trainers and training courses. It is essential that there should be minimal adverse effects on trainees in radiology and sonography. It must be recognised that training requires additional time, space and equipment. Training should be properly costed and funded.
- 2.12 Training should be related to the specialist requirements of the trainee; that is, training should be modular. Within any one level of training, it may be appropriate for a trainee to become proficient in some but not all of the individual modules and only undertake ultrasound practice in this/these areas.
- 2.13 Training should be given in departments which have a multidisciplinary (medical, surgical, radiological and so on) philosophy, an adequate throughput of work, a radiologist or Level 2/3 sonographer practitioner with experience and an interest in training in the module required, appropriate equipment and an active audit process. The role of sonographer practitioners in delivering some or all of this modular training should be formally recognised and agreed.
- 2.14 Regular appraisal should take place during the training period. It must be recognised that not all trainees have the aptitude to undertake ultrasound scanning and that, some, despite undergoing training, may not acquire the appropriate skills ever to practise independently. At the end of a period of training, a 'competency assessment' form should be completed for each trainee, which will determine the area or

areas in which they can practise independently (see Section 3). The responsibility to be adequately trained and to maintain those skills lies with the individual practising ultrasound. An assessment of competence is a reflection on the position at the time the assessment is undertaken and no more. If sonographic practitioners are involved with competence assessment then they should be fully supported in this respect by a responsible radiologist experienced in ultrasound or another ultrasound Level 2/3 trained medical practitioner.

- 2.15 Following training, regular and relevant continuing professional development (CPD) should be undertaken and documented. It is the responsibility of the trainee to ensure that their practical skills are maintained by ensuring that regular ultrasound sessions are undertaken and that there is an adequate range of pathology seen in their ultrasound practice.

# 3 Training recommendations

Training should consist of both theoretical and practical syllabuses.

## 3.1 Theoretical training

3.1.1 Preliminary theoretical training should cover the physics of ultrasound, levels and sophistication of equipment, image recording, reporting, artefacts and the relevance of other imaging modalities to ultrasound. This element of training may be best delivered by linking with some of the excellent courses run by university departments accredited by the Consortium for the Accreditation of Sonographic Education (CASE).

The Radiology-integrated Training Initiative (R-ITI), a free resource, includes eight sections on theoretical principles.

3.1.2 The theoretical syllabus is set out in Appendix 11.

## 3.2 Practical training

3.2.1 A syllabus for each area of ultrasound specialisation structured into the three levels of training has been developed, incorporating theoretical training on anatomy and pathology and a practical syllabus listing conditions which should be included in the experience of the trainee (Appendices 1–9). A modular anatomical approach is recommended for musculoskeletal ultrasound (for example, a trainee may become proficient in shoulder ultrasound alone), as set out in Appendix 10. In other areas of ultrasound specialisation, in appropriate circumstances, a limited anatomical or modular approach may also be acceptable if full competence in that area is demonstrated and future clinical practice is confined to that area alone.

3.2.2 Practical experience should be gained under the guidance of a named supervisor trained in ultrasound within a training department. In the context of advice from the RCR, this would normally be in a department of clinical radiology. There may be some areas of ultrasound practice which are not covered by these modules such as intra-operative ultrasound and transcranial Doppler ultrasound. Where required, training modules based on similar principles should be developed for any area of ultrasound practice not covered in this publication.

3.2.3 The syllabuses set out in Appendices 1–10 include a competency assessment sheet for training. This should be completed during the course of training as it will help to determine in which areas(s) the trainee can practise independently (see Section 2.14).

3.2.4 The requirements for the different levels of training are as follows:

### 3.2.4.1 Level 1

Different trainees will acquire the necessary skills at different rates and the endpoint of the training programme should be judged by an assessment of practical competence.

Examinations should encompass the full range of pathological conditions listed in the syllabuses.

A logbook listing the number and type of examinations undertaken by the trainee themselves should be kept.

An illustrated logbook of specific normal and abnormal findings may be appropriate for some syllabuses.

Training should usually be supervised by a Level 2/3 practitioner. In certain circumstances it may be appropriate to delegate some or most of this supervision to a Level 1 practitioner with at least two years' experience of Level 1 practice.

### 3.2.4.2 Level 2

This usually requires at least one year of experience at Level 1, with the equivalent of at least one session per week.

A significant further number of examinations should have been undertaken in order to encompass the full range of conditions and procedures encountered in each module.

A logbook listing the numbers and types of examinations undertaken by the trainee should be maintained.

An illustrated logbook of specific normal and abnormal findings may be appropriate for some syllabuses.

Supervision of training should be undertaken by someone who has achieved at least Level 2 competence and has had at least two years' experience at that Level.

#### *3.2.4.3 Level 3*

This requires practitioners to spend a significant part of their time undertaking ultrasound examinations, teaching, research and development.

They will have spent a continuous period of sub-specialist training in which ultrasound will have been a significant component.

They will be able to perform specialised examinations at the leading edge of ultrasound practice such as the use of intravascular ultrasound contrast agents and the performance of advanced ultrasound-guided invasive procedures.

3.2.5 The syllabuses for each area of ultrasound specialisation are outlined in Appendices 1–10.

# 4 Continuing professional development

- 4.1 The minimum amount of ongoing experience in ultrasound as outlined in each syllabus should be maintained.
- 4.2 CPD should be undertaken which incorporates elements of ultrasound practice. This should be included in the annual appraisal and revalidation process.
- 4.3 Regular audit of the individual's ultrasound practice should be undertaken to demonstrate that the indications, performance and diagnostic quality of the service are all satisfactory. The audit process should be independent and the format should be in line with RCR recommendations. Evidence of audit should be available to commissioners of service if required. Audit templates can be submitted via the RCR's Clinical Radiology AuditLive.
- 4.4 The individual should keep up to date with the relevant literature.
- 4.5 The individual should attend regular multidisciplinary meetings and have an ultrasound mentor.

Approved by the Board of the Faculty of Clinical Radiology: 15 June 2012

# References

1. The Royal College of Radiologists. *Focused ultrasound training standards*. London: The Royal College of Radiologists, 2012.
2. European Federation of Societies for Ultrasound in Medicine and Biology (EFSUMB). *Minimum training requirements for the practice of medical ultrasound in Europe*. [www.efsumb.org/guidelines/2009-04-14apx1.pdf](http://www.efsumb.org/guidelines/2009-04-14apx1.pdf) (last accessed 23/11/12)

# Appendix 1. Urological ultrasound

This curriculum is intended for clinicians who perform diagnostic urological ultrasound and ultrasound-guided urological intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

## Focused training and practice

There are frequent situations arising in clinical practice where rapid bedside assessment using focused ultrasound techniques can help with the assessment of, and treatment planning for, patients. In this situation, rapid ultrasound assessment by a competent non-radiological clinician may be more appropriate than waiting for a formal ultrasound list during normal working hours. Some clinicians may wish to focus on only one skill such as deciding whether the collecting systems/ureters are dilated or not, placing a suprapubic catheter, transrectal ultrasound (TRUS) biopsies and so on. These unitary skills may aid the clinician's practice and greatly improve patient pathways.

Each clinician will have their own requirement for focused training and to accommodate their training requirements, a targeted curriculum and syllabus should be created by local trainers, drawing on appropriate elements of the knowledge base and competencies to be acquired from Levels 1–2, depending on the level of practice expected. An example syllabus is included in the focused ultrasound document; for example, suprapubic catheterisation.<sup>1</sup>

It is essential that all ultrasound examinations that may have any influence on patient management are performed by individuals who are competent to provide an accurate examination and assessment and that the images and a formal report are recorded on a RIS/PACS system.

## Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasonic anatomy
  - Kidneys
  - Ureters
  - Other retroperitoneal structures (adrenals, aorta, inferior vena cava [IVC])
  - Bladder
  - Seminal vesicles
  - Prostate
  - Scrotal contents
  - Other pelvic structures (uterus, ovaries, lymph nodes, vessels, bowel)
- Pathology in relation to ultrasound
  - Kidneys: congenital anomalies, cysts, tumours (benign and malignant), stones, collecting system dilatation, renal and peri-renal abscesses, trauma, diffuse renal diseases
  - Ureters: dilatation, obstruction
  - Bladder: tumours, diverticula, wall thickening, calculi, volume estimation
  - Prostate: infection, hyperplasia, tumours
  - Scrotal contents: testicular tumours, cysts, torsion, hydrocele, inflammatory problems, trauma

## Level 1. Training and practice

Practical training should involve at least one ultrasound list per week over a period of 3–6 months, with approximately 5–10 examinations performed by the trainee (under supervision) per session.

A minimum of 250 examinations should be undertaken. However, different trainees will acquire the necessary skills at different rates, and the endpoint of the training programme should be judged by an assessment of competencies.

Examinations should encompass the full range of pathological conditions listed below.

A logbook listing the types of examinations undertaken should be kept.

Training should be supervised either by someone who has obtained at least Level 2 competence in urological ultrasound or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should attend an appropriate theoretical course and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

## Level 1. Competencies to be acquired

### *Kidneys*

To be able to:

- Perform a thorough ultrasound examination of the kidneys in different planes
- Recognise normal renal ultrasonic anatomy and common normal variants
- Measure renal length and assess variation from normality
- Recognise and assess the degree of collecting system dilatation
- Recognise and diagnose simple cysts
- Recognise complex cysts and refer for appropriate further investigation
- Recognise renal tumours and refer for appropriate further investigation
- Recognise and diagnose renal stones
- Recognise peri-renal abnormalities and refer for appropriate further investigation
- Recognise abnormalities which need referral for scanning by a more experienced Ultrasonologist and/or further investigation

### *Bladder*

To be able to:

- Perform a thorough ultrasound examination of the bladder in different planes
- Recognise normal ultrasonic anatomy of the bladder and common normal variants
- Measure bladder volume
- Recognise and diagnose bladder diverticula
- Recognise and assess bladder tumours
- Recognise bladder calculi
- Use colour Doppler to assess ureteric jets
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Scrotum*

To be able to:

- Perform a thorough ultrasound examination of the scrotal contents in different planes
- Recognise normal ultrasonic anatomy of the testes and epididymi and common normal variants
- Recognise and diagnose epididymal cysts
- Recognise and diagnose varicoceles
- Use Doppler to help differentiate torsion/inflammatory problems
- Recognise and assess intra-scrotal and intra-testicular calcifications
- Recognise and assess testicular tumours
- Recognise inflammatory changes in testes and epididymides
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Prostate*

To be able to:

- Recognise normal ultrasonic anatomy and common normal variants
- Perform transrectal ultrasound
- Measure prostatic volume
- Identify abnormal focal lesions
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Other*

- To be able to recognise and, where appropriate, refer for further investigation:
  - Normal aorta and aortic aneurysm
  - Normal liver and liver masses
  - Normal uterus and ovaries and gynaecological masses
- To be able to use ultrasound in the assessment of patients presenting with:
  - Haematuria
  - Loin pain/renal colic

- Loin mass
- Lower urinary tract symptoms
- Recurrent urinary tract infections
- Suprapubic mass
- Palpable masses in the scrotum
- Scrotal pain

## **Level 2. Knowledge base**

- Physics and technology
  - In-depth knowledge and understanding of the physics of ultrasound
  - In-depth knowledge and understanding of the technology of ultrasound equipment
- Ultrasound techniques
  - The advanced use of Doppler ultrasound, including spectral, colour and power Doppler
  - The use of ultrasound for guiding interventional procedures
  - Further applications of transabdominal ultrasound
  - Further application of endocavity ultrasound (eg, transvaginal and transrectal ultrasound)
  - Intraoperative ultrasound
- Sectional and ultrasonic anatomy
  - The normal renal and pelvic vasculature, including an understanding of the Doppler signals obtained from these vessels
  - More detailed knowledge of structures outside the urinary tract in the abdomen and pelvis
  - Ultrasound anatomy of the penis and female genital organs

## **Level 2. Training and practice**

Practical training should involve at least one year of experience at Level 1 with a minimum of one session per week.

A further 600 examinations should have been undertaken in order to encompass the full range of conditions and procedures referred to below.

A logbook listing all examinations undertaken should be kept.

Supervision of training should be undertaken by someone who has achieved at least Level 2 competence in urological ultrasound, has had at least two years' experience at that Level, and who would normally be of consultant status.

A Level 2 practitioner will be able to accept referrals from Level 1 practitioners.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice, and refined during a period of clinical practice.

*Kidneys, bladder, prostate, scrotal contents*

To be able to:

- Recognise all pathology affecting the urinary tract and provide an accurate diagnosis in almost all cases
- Recognise abnormalities which are out with their experience and refer on appropriately
- Perform ultrasound-guided invasive procedures, including cyst aspiration, abscess drainage, renal biopsy, percutaneous nephrostomy, suprapubic bladder catheter insertion and transrectal prostate biopsies
- Perform Doppler ultrasound studies relevant to the urinary tract
- Recognise abnormalities elsewhere in the abdomen and pelvis which need referral for scanning by another ultrasonologist and/or further investigation

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking urological ultrasound, teaching, research and development and will be an 'expert' in this area.

They will have spent a continuous period of specialist training in urological ultrasound.

They will perform specialised examinations at the leading edge of ultrasound practice.

They will accept tertiary referrals from Level 1 and Level 2 practitioners and will perform specialised examinations (eg, the use of intravascular ultrasound agents in evaluating possible malignancy) as well as performing advanced ultrasound-guided invasive procedures.

### **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

A specialist registrar will need to continue to perform ultrasound scans throughout the remainder of their training programme. Such further ultrasound practice may be intermittent, but no more than three months should elapse without trainees using their ultrasound skills, and at least 100 examinations should be performed per year.

A medical practitioner performing Level 1 ultrasound should continue to perform at least 250 ultrasound examinations per year on a regular basis, should have regular meetings with radiological colleagues and should have a named radiologist as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

Approved by the British Association of Urological Surgeons.

# Appendix 1. Urological ultrasound training competency assessment sheet

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
Use of the local RIS and PACS system					
Competencies/skills to be acquired — Level 1					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Kidneys</b>			<b>Scrotum</b>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Ultrasonic anatomy and common normal variants</li> <li>▪ Renal length and variation from normality</li> <li>▪ Collecting system dilatation</li> <li>▪ Simple cysts</li> <li>▪ Complex cysts</li> <li>▪ Tumours</li> <li>▪ Stones</li> </ul>			<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Ultrasonic anatomy and common normal variants</li> <li>▪ Epididymal cysts</li> <li>▪ Varicoceles</li> <li>▪ Intrascrotal and intratesticular calcifications</li> <li>▪ Tumours</li> <li>▪ Inflammatory changes in testes and epididymides</li> <li>▪ Use Doppler to help differentiate torsion/inflammatory problems</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Peri-renal abnormalities</li> </ul>			<b>Other</b>		
<b>Bladder</b>			<ul style="list-style-type: none"> <li>– Normal aorta and aortic aneurysm</li> <li>– Normal liver and liver masses</li> <li>– Normal uterus and ovaries and gynaecological masses</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Ultrasonic anatomy and common normal variants</li> </ul>			<ul style="list-style-type: none"> <li>▪ Use ultrasound in the assessment of patients presenting with:</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Bladder volume</li> </ul>			<ul style="list-style-type: none"> <li>– Haematuria</li> <li>– Loin pain/renal colic</li> <li>– Loin mass</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Diverticula</li> <li>▪ Tumours</li> <li>▪ Calculi</li> <li>▪ Use colour Doppler to assess ureteric jets</li> </ul>			<ul style="list-style-type: none"> <li>– Lower urinary tract symptoms</li> <li>– Recurrent urinary tract infection</li> </ul>		
<b>Prostate</b>			<ul style="list-style-type: none"> <li>– Suprapubic mass</li> <li>– Palpable scrotal masses</li> <li>– Scrotal pain</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Ultrasonic anatomy and common normal variants</li> <li>▪ Transrectal ultrasound</li> <li>▪ Prostatic volume</li> <li>▪ Abnormal focal lesions</li> </ul>			<ul style="list-style-type: none"> <li>▪ Know when to refer to a more expert ultrasonologist</li> </ul>		
Core knowledge base — Level 2		Trainee		Trainer	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Sectional and ultrasonic anatomy		
Ultrasound techniques					

Competencies/skills to be acquired — Level 2					
To be competent to perform/recognise the following:					
Pathology affecting the urinary tract and provide an accurate diagnosis in almost all cases			Doppler ultrasound studies relevant to the urinary tract		
Abnormalities which are out with personal experience and refer on appropriately			Abnormalities elsewhere in the abdomen and pelvis which need referral for scanning by another ultrasonologist and/or further investigation		
Ultrasound-guided invasive procedures, including cyst aspiration, abscess drainage, renal biopsy, percutaneous nephrostomy trans-rectal prostate biopsies and suprapubic bladder catheter insertion					

## Appendix 2. Gynaecological ultrasound

This curriculum is intended for clinicians who perform diagnostic ultrasound and ultrasound-guided intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

The aim is to provide high-quality diagnostic ultrasound for females.

Members of the RCR and RCOG who are in training will fulfil the requirements of these Colleges.

The training outlined below is provided for the benefits of safe practice for medical practitioners who wish to provide diagnostic ultrasound for females such as GPs.

### Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration and report writing (see Appendix 11)
- The techniques of transabdominal and transvaginal scanning are essential.
- A full understanding of the issues relating to the performance of intimate examinations and the importance of informed consent for the procedures is emphasised.
- Sectional and ultrasonic anatomy
  - Uterus (including physiological changes with age and cycle)
  - Ovaries (including physiological changes with age and menstrual cycle)
  - Cervix and vagina
  - First trimester gestation appearances
  - Bladder and urethra
  - Associated structures; omentum and peritoneal fluid
- Pathology in relation to ultrasound
  - Uterus: fibroids, adenomyosis, intrauterine contraceptive devices (IUCDs), endometrial hyperplasia, polyps and tumours
  - Ovaries: cysts and their complications, endometrioma, tumours, inflammation and infection, polycystic and hyperstimulated ovaries, torsion
  - Fallopian tubes: hydro/pyo-salpinges
  - Cervix and vagina: congenital lesions, cysts, tumour, retained foreign bodies
  - First trimester: location, viability, biometry, to include fetal number and chorionicity, ectopic pregnancy; signs of non-viability; haemorrhage; retained products of conception
  - Bladder and urethra: volume estimation diverticula, wall thickening, calculi, tumours, peri-urethral cysts and abscesses
  - Other pelvic pathology to recognise deviation from normal eg free fluid or masses

### Level 1. Training and practice

Practical training should involve at least 30 ultrasound sessions within a period of six months with approximately 3–8 examinations performed by the trainee (under supervision) per session. However, different trainees will acquire the necessary skills at different rates and the endpoint of the training programme should be judged by an assessment of competencies to perform and report an ultrasound examination.

Examinations should ideally encompass the full range of pathological conditions listed below.

For some practitioners where Level 1 competencies in only one area of practice the training and competency assessment in these areas only can be obtained (eg, in the assessment of early pregnancy clinics or postmenopausal patients).

A logbook listing the type of examinations undertaken should be kept.

Training should be supervised either by someone who has obtained at least Level 2 competence in gynaecological ultrasound, or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should attend an appropriate theoretical course and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

## Level 1. Competencies to be acquired

### *Early pregnancy*

To be able to:

- Define pregnancy locality by ultrasound signs
- Recognise signs of viability/non-viability
- Recognise normal appearances as gestation advances
- Recognise signs of ectopic pregnancy
- Identify multiple pregnancy and chorionicity
- Date pregnancy by crown rump length (CRL)
- Recognise signs of haemorrhage
- Recognise signs of retained products of conception
- Understand the role of ultrasound in the setting of early pregnancy clinical pathways and laboratory findings
- Understand the terms and recognise the ultrasound findings in pregnancies of unknown location and pregnancy of uncertain viability
- Write a structured report of the ultrasound findings

### *Abnormal vaginal bleeding*

To be able to:

- Recognise normal and abnormal endometrial thickness
- Recognise the features of endometrial polyps/carcinoma
- Recognise the features of atrophic endometrium
- Recognise features of fibroids and their localisation
- Recognise appearances of IUCDs and their localisation
- Recognise when further investigation is required and what to ask for
- Understand the need for further referral and clinical pathways
- Write a structured report of the ultrasound findings

### *Pelvic pain*

To be able to:

- Recognise the features of ovarian cyst torsion, rupture or haemorrhage
- Recognise the features of endometrioma
- Recognise appearances of hydrosalpinges
- Recognise the features of pelvic inflammatory disease
- Recognise non-gynaecological causes of pelvic pain
- Understand the need for further referral and clinical pathways
- Write a structured report of the ultrasound findings

### *Pelvic mass*

To be able to:

- Recognise typical appearances of uterine and ovarian masses
- Recognise features suggesting benign or malignant pathology
- Variations from normal suggesting non-gynaecological causes of a pelvic mass
- Understand the need for further referral and clinical pathways
- Write a structured report of the ultrasound findings

### *Reproductive medicine*

To be able to:

- Recognise features of the endometrium at different stages of the menstrual cycle
- Recognise features of the ovary at different stages of the menstrual cycle
- Recognise features of a stimulated and hyperstimulated ovary
- Recognise features of a polycystic ovary
- Understand the need for further referral and clinical pathways
- Write a structured report of the ultrasound findings

### *Bladder*

To be able to:

- Perform an ultrasound examination of the bladder in different planes
- Recognise normal anatomy of the bladder and common normal variants
- Measure bladder volume
- Recognise and diagnose bladder diverticula
- Recognise bladder tumours
- Recognise bladder calculi
- Recognise variations from normal/abnormalities which need referral for scanning by a more experienced ultrasonologist and/or further investigation
- Understand the need for further referral and clinical pathways
- Write a structured report of the ultrasound findings

Practitioners should:

- Include ultrasound in their ongoing CPD
- Audit their practice

## **Level 2. Knowledge base**

- Physics and technology
  - In-depth knowledge and understanding of the physics of ultrasound
  - In-depth knowledge and understanding of the technology of ultrasound equipment
- Ultrasound techniques
  - The advanced use of Doppler ultrasound, including spectral, colour and power Doppler
  - The use of ultrasound for guiding interventional procedures
  - Further applications of transabdominal ultrasound
  - Further applications of transvaginal ultrasound: saline infusion hystero-graphy (SIH), HyCoSy
- Sectional and ultrasonic anatomy
  - The normal pelvic and gynaecological organ vasculature, including an understanding of the Doppler signals obtained from these vessels
  - More detailed knowledge of structures outside the female reproductive tract in the pelvis

## **Level 2. Training and practice**

Practical training should include at least one year of experience at Level 1 with a minimum of the equivalent of one session per week.

A further 600 examinations should have been undertaken in order to encompass the full range of conditions and procedures referred to below.

A logbook listing all examinations undertaken should be kept.

Supervision of training should be undertaken by someone who has achieved at least Level 2 competence in gynaecological ultrasound, has had at least two years' experience at that level, and who would normally be of consultant status.

A Level 2 practitioner will be able to accept referrals from Level 1 practitioners.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice, and refined during a period of clinical practice.

### *Female reproductive tract*

To be able to:

- Recognise and correctly diagnose almost all pathology affecting the female genital tract
- Perform Doppler ultrasound studies relevant to the uterus and ovaries
- Recognise abnormalities elsewhere in the and pelvis which need referral for scanning by another ultrasonologist and/or further investigation

In addition specifically to be able to recognise and evaluate:

- Causes of an abnormal Doppler waveform
- Changes associated with precocious puberty, thelarche, adrenarche and virilisation
- Congenital anomalies
- Features of lymph nodes in the inguinal and iliac chains
- Bartholin's cysts, abscesses and periurethral lesions

- Features of haematocolpos
- Features of adenomyosis
- Non-ovarian endometriosis
- Non-gynaecological causes of pelvic pain and how to diagnose appendicitis, inflammatory bowel disease, bowel cancer, hernias, aneurysms and bladder disease
- Different types of complex ovarian masses
- The principles of oocyte collection by transvaginal ultrasound-guided aspiration of follicles

Practitioners should:

- Include ultrasound in their ongoing CPD
- Audit their practice.

### **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking gynaecological ultrasound, teaching, research and development.

They will have spent a continuous period of specialist training in gynaecological ultrasound.

They will accept tertiary referrals from Level 1 and 2 practitioners.

They will perform specialised examinations at the leading edge of ultrasound practice.

#### **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

Once trained and assessment of competencies confirmed, the practitioner will need to continue to perform ultrasound throughout the remainder of their training programme. Such further ultrasound practice may be intermittent, but no more than three months should elapse without the trainee using their scanning skills, and competency with evidence of sufficient scanning to maintain these competencies.

All practitioners should work in a team in order to maintain competencies.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

## Appendix 2. Gynaecological ultrasound competency sheet

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
Techniques of transabdominal and transvaginal scanning			Issues regarding intimate examinations and informed consent		
<b>Competencies/skills to be acquired — Level 1</b>					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Early pregnancy</b>			<b>Reproductive medicine</b>		
▪ Dating pregnancy			▪ Endometrial changes with menstrual cycle		
▪ Signs of non-viability			▪ Ovarian changes with menstrual cycle		
▪ Multiple pregnancy and chorionicity			▪ Polycystic ovaries		
▪ Haemorrhage			▪ Stimulated and hyperstimulated ovaries		
▪ Retained products of conception			<b>Kidneys</b>		
▪ Ectopic pregnancy			▪ Ultrasound examination in different scan planes		
<b>Vaginal bleeding</b>			▪ Ultrasonic anatomy and common normal variants		
▪ Fibroids and their localisation			▪ Renal length		
▪ IUCDs and their localisation			▪ Collecting system dilatation		
▪ Normal and abnormal endometrial thickness			▪ Cysts		
▪ Atrophic and hyperplastic endometrium			▪ Tumours		
▪ Endometrial polyps			▪ Stones		
<b>Pelvic pain</b>			<b>Bladder</b>		
▪ Ovarian cyst complications			▪ Ultrasound examination in different planes		
▪ Endometriosis			▪ Ultrasonic anatomy and common normal variants		
▪ Pelvic inflammatory disease			▪ Bladder volume		
▪ Non-gynaecological causes of pain			▪ Bladder diverticula		
<b>Pelvic mass</b>			▪ Tumours		
▪ Identification			▪ Calculi		
▪ Organ of origin of mass			▪ Ultrasound examination in different planes		
▪ Benign versus malignant features			▪ Ultrasonic anatomy and common normal variants		
			<b>General</b>		
			▪ Know when to refer to a more expert ultrasonologist		
<b>Core knowledge base — Level 2</b>		<b>Trainee</b>		<b>Trainer</b>	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Sectional and ultrasonic anatomy		
Ultrasound techniques					

Competencies/skills to be acquired — Level 2					
To be competent to perform/recognise/diagnose/evaluate the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Female reproductive tract</b>					
<ul style="list-style-type: none"> <li>▪ Almost all pathology affecting the female genitourinary tract</li> </ul>			<ul style="list-style-type: none"> <li>▪ Bartholin's cysts, abscesses and periurethral lesions</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Ultrasound-guided invasive procedures, including ascitic drainage, omental biopsy, pelvic mass biopsy (transabdominal or transvaginal), lymph node aspiration, SIH and HyCoSy</li> </ul>			<ul style="list-style-type: none"> <li>▪ Features of haematocolpos</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Doppler ultrasound studies relevant to the uterus and ovaries</li> </ul>			<ul style="list-style-type: none"> <li>▪ Features of adenomyosis</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Abnormalities elsewhere in the abdomen and pelvis which need</li> </ul>			<ul style="list-style-type: none"> <li>▪ Non-ovarian endometriosis</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Referral for scanning by another ultrasonologist and/or further investigation</li> </ul>			<ul style="list-style-type: none"> <li>▪ Non-gynaecological causes of pelvic pain and how to diagnose appendicitis, inflammatory bowel disease, bowel cancer, hernias, aneurysms and bladder disease</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Stage ovarian and uterine tumours</li> </ul>			<ul style="list-style-type: none"> <li>▪ Different types of complex ovarian masses</li> </ul>		
<b>Other</b>			<ul style="list-style-type: none"> <li>▪ Malignant disease of the omentum, peritoneum and the rest of the abdomen</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Causes of an abnormal Doppler waveform</li> </ul>			<ul style="list-style-type: none"> <li>▪ Features of pleural effusions</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Changes associated with precocious puberty, thelarche, adrenarache and virilisation</li> </ul>			<ul style="list-style-type: none"> <li>▪ Common sites and features of tumours that metastasise to the pelvis</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Congenital anomalies</li> </ul>			<ul style="list-style-type: none"> <li>▪ The principles of oocyte collection by transvaginal ultrasound-guided aspiration of follicles</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Features of lymph nodes in the inguinal and iliac chains</li> </ul>					

## Appendix 3. Gastrointestinal ultrasound

This curriculum is intended for clinicians who perform diagnostic gastrointestinal ultrasound and ultrasound-guided intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

### Focused training and practice

There are frequent situations arising in clinical practice where rapid bedside assessment using focused ultrasound techniques can help with the assessment of, and treatment planning for, patients. In this situation, rapid ultrasound assessment by a competent non-radiological clinician may be more appropriate than waiting for a formal ultrasound list during normal working hours. Some clinicians may wish to focus on only one skill such as drainage of ascites and ultrasound-guided liver biopsies. These unitary skills may aid the clinician's practice and greatly improve patient pathways.

Each clinician will have their own requirement for focused training and to accommodate their training requirements, a targeted curriculum and syllabus should be created by local trainers, drawing on appropriate elements of the knowledge base and competencies to be acquired from Levels 1–2, depending on the level of practice expected. An example syllabus is included in the focused ultrasound document; for example, drainage of ascites and ultrasound-guided liver biopsies.<sup>1</sup>

It is essential that all ultrasound examinations that may have any influence on patient management are performed by individuals who are competent to provide an accurate examination and assessment and that the images and a formal report are recorded on a RIS/PACS system.

### Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasonic anatomy
  - Liver
  - Gallbladder
  - Bile ducts
  - Pancreas
  - Spleen
  - Kidneys
  - Other structures (uterus, ovaries, lymph nodes, vessels, bowel)
- Pathology in relation to ultrasound
  - Liver: cysts, benign and malignant tumours, metastatic disease, fatty change, cirrhosis
  - Biliary system: gallbladder stones, acute and chronic cholecystitis, gallbladder tumours, bile duct obstruction including level of obstruction, intrahepatic duct gas and stones
  - Pancreas: pancreatitis, duct stones, duct dilatation, pancreatic tumours
  - Portal venous system and spleen: splenic enlargement, portal venous distention, varices, thrombosis, ascites
  - Other structures: renal masses and urinary tract obstruction (hydronephrosis), ovarian and uterine masses including cysts, tumours, fibroids and unexpected pregnancy

### Level 1. Training and practice

Practical training should involve at least one ultrasound list per week over 3–6 months with approximately 5–10 examinations performed by the trainee (under supervision) per session. A minimum of 250 examinations should be undertaken. However, different trainees will acquire the necessary skills at different rates and the endpoint of the training programme should be judged by an assessment of competencies.

Examinations should encompass the full range of pathological conditions listed below.

A logbook listing the types of examinations undertaken should be kept.

Training should be supervised either by someone who has obtained at least Level 2 competence in gastrointestinal ultrasound or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should attend an appropriate theoretical course and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

## **Level 1. Competencies to be acquired**

### *Liver*

To be able to:

- Perform a thorough ultrasound examination of the liver in different scan planes
- Recognise normal hepatic anatomy and variants
- Recognise normal and abnormal liver texture, such as fatty change, cirrhosis, atrophy and hypertrophy
- Recognise focal lesions, particularly cysts and benign lesions, and be able to detect possible malignant lesions requiring further investigation
- Recognise normal hepatic and portal venous anatomy within the liver
- Recognise abnormalities of the hepatic and portal venous system, eg, thrombosis
- Perform ultrasound-controlled biopsy for the evaluation of parenchymal liver disease

### *Biliary system*

To be able to:

- Perform a thorough evaluation of the biliary system
- Recognise normal ultrasonic anatomy of the biliary system and its frequent normal variants
- Recognise abnormalities of the gallbladder wall
- Recognise gallbladder stones
- Recognise features of acute and chronic cholecystitis
- Assess bile duct dilatation at intrahepatic and extrahepatic levels and determine level of obstruction

### *Pancreas*

To be able to:

- Perform a thorough examination of the pancreas recognising normal anatomy
- Recognise the limitations of pancreatic ultrasound because of bowel gas
- Recognise solid and cystic tumours within the head and body of the pancreas
- Recognise pancreatic duct dilatation and pancreatic duct stones
- Recognise the features of acute and chronic pancreatitis and their complications

### *Portal venous system and spleen*

To be able to:

- Evaluate the size of the spleen and recognise focal lesions and evidence of trauma
- Evaluate the portal vein and its diameter and the presence of portal venous thrombosis
- Recognise cavernous transformation of the portal vein and varices

### *Bowel*

To be able to:

- Recognise normal stomach, small and large bowel
- Recognise focal intestinal abnormality and understand the principles of further investigation
- Recognise small bowel obstruction

### *Other*

To be able to:

- Recognise abdominal aortic aneurysm
- Recognise hydronephrosis
- Recognise normal kidneys, uterus and ovaries
- Recognise renal and gynaecological masses

## **Level 2. Knowledge base**

### *Sectional and ultrasonic anatomy*

- Detailed understanding of intestinal, mesenteric, peritoneal, omental, vascular and retroperitoneal anatomy

### *Pathology in relationship to ultrasound*

- An understanding of disease processes which affect the peritoneal cavity, its mesenteries, ligaments and compartments
- An understanding of the pathways of spread of intraperitoneal and retroperitoneal disease

## **Level 2. Training and practice**

Practical training should involve at least one year of experience at Level 1 with a minimum of one session per week.

A further 500 examinations should have been undertaken in order to encompass the full range of conditions and procedures listed below.

A logbook listing all examinations undertaken should be kept.

Supervision of training should be undertaken by someone who has achieved Level 2 competence in gastrointestinal ultrasound, has had at least two years' experience at that level and who would normally be of consultant status.

A Level 2 practitioner will be able to accept referrals from Level 1 practitioners.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice and refined during a period of clinical practice.

- Perform a comprehensive ultrasound examination of all of the solid organs within the abdomen.
- Be able to evaluate the small bowel for focal or diffuse disease.
- Be able to evaluate the bowel in inflammatory bowel disease.
- Be able to evaluate the large bowel for the presence of diverticular disease, appendicitis and their complications, tumours and obstruction.
- Be able to evaluate the peritoneal cavity, its mesenteries, compartments and the omentum for the presence of infective or malignant disease.
- Be able to undertake ultrasound-guided drainage of peritoneal fluid collections.
- Be able to evaluate the hepatic and portal venous systems using spectral, colour and power Doppler ultrasound.
- Be able to undertake ultrasound-guided biopsy of focal liver lesions.

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking gastrointestinal ultrasound, teaching, research and development and will be an 'expert' in this area.

They will accept tertiary referrals from Level 1 and 2 practitioners and will perform specialised examinations (eg, the use of intravascular ultrasound agents in evaluating focal liver lesions) as well as performing advanced ultrasound-guided invasive procedures.

## **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

A specialist registrar will need to continue to perform ultrasound throughout the remainder of their training programme. Such further ultrasound practice may be intermittent, but no more than three months should elapse without the trainee using their scanning skills, and at least 100 scans should be performed per year.

A medical practitioner performing ultrasound at Level 1 should continue to perform at least 250 examinations per year on a regular basis, should have regular meetings with radiological colleagues and should have a named radiologist as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

## Appendix 3 competency sheet: gastrointestinal ultrasound

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
<b>Competencies/skills to be acquired — Level 1</b>					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Liver</b>			<b>Pancreas</b>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different scan planes</li> <li>▪ Normal hepatic anatomy and variants</li> <li>▪ Normal and abnormal liver texture</li> <li>▪ Atrophy and hypertrophy</li> <li>▪ Fatty change</li> <li>▪ Cirrhosis</li> <li>▪ Cysts and benign lesions</li> <li>▪ Malignant lesions</li> <li>▪ Normal hepatic and portal venous anatomy</li> <li>▪ Abnormalities of the portal venous system eg, thrombosis</li> <li>▪ Ultrasound-controlled biopsy for the evaluation of parenchymal liver disease</li> </ul>			<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different scan planes</li> <li>▪ Normal anatomy and variants</li> <li>▪ Solid and cystic tumours</li> <li>▪ Duct dilatation and stones</li> <li>▪ Acute and chronic pancreatitis and complications</li> </ul>		
			<b>Portal venous system and spleen</b>		
			<ul style="list-style-type: none"> <li>▪ Splenic size and focal lesions</li> <li>▪ Splenic trauma</li> <li>▪ Portal vein diameter</li> <li>▪ Cavernous transformation of the portal vein</li> </ul>		
			<ul style="list-style-type: none"> <li>▪ Varices</li> </ul>		
<b>Biliary system</b>			<b>Other</b>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different scan planes</li> <li>▪ Normal ultrasonic anatomy and variants</li> <li>▪ Abnormalities of the gallbladder wall including gall bladder tumours</li> <li>▪ Gallbladder stones</li> <li>▪ Acute and chronic cholecystitis</li> <li>▪ Bile duct dilatation and level of obstruction</li> </ul>			<ul style="list-style-type: none"> <li>▪ Normal uterus, ovaries, kidneys and aorta</li> <li>▪ Abdominal aortic aneurysm</li> <li>▪ Hydronephrosis</li> </ul>		
			<ul style="list-style-type: none"> <li>▪ Renal and gynaecological masses</li> </ul>		
			<b>General</b>		
			<ul style="list-style-type: none"> <li>▪ Know when to refer to a more expert ultrasonologist</li> </ul>		
<b>Bowel</b>					
<ul style="list-style-type: none"> <li>▪ Normal stomach, small and large bowel</li> <li>▪ Focal intestinal abnormality</li> <li>▪ Small bowel obstruction</li> </ul>					
<b>Competencies/skills to be acquired — Level 2</b>					
To be competent to perform/ diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
Ultrasound examination of all of the solid organs within the abdomen			Bowel tumour		
Focal or diffuse small bowel disease			Infective or malignant disease of peritoneum, mesenteries and omentum		

Inflammatory bowel disease			Ultrasound-guided drainage of peritoneal fluid collections		
Diverticular disease, appendicitis and their complications			The hepatic and portal venous systems using spectral, colour and power Doppler ultrasound		
Bowel obstruction			Ultrasound-guided biopsy of focal liver lesions		

## Appendix 4. Vascular ultrasound

This curriculum is intended for clinicians who perform diagnostic vascular ultrasound and ultrasound-guided intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

### Focused training and practice

There are frequent situations arising in clinical practice where rapid bedside assessment using focused ultrasound techniques can help with the assessment of, and treatment planning for, patients. In this situation, rapid ultrasound assessment by a competent non-radiological clinician may be more appropriate than waiting for a formal ultrasound list during normal working hours. Some clinicians may wish to focus on only one skill such as to provide ultrasound-guided vascular cannulation. These unitary skills may aid the clinician's practice and greatly improve patient pathways.

Each clinician will have their own requirement for focused training and in order to accommodate their training requirements, a targeted curriculum and syllabus should be created by local trainers, drawing on appropriate elements of the knowledge base and competencies to be acquired from Levels 1–2, depending on the level of practice expected. An example syllabus is included in the focused ultrasound document; for example, ultrasound-guided vascular cannulation.<sup>1</sup>

It is essential that ALL ultrasound examinations that may have any influence on patient management are performed by individuals who are competent to provide an accurate examination and assessment and that the images and a formal report are recorded on a RIS/PACS system.

### Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- To have full working knowledge of the principles, techniques, instrumentation and practical working knowledge of real-time and Doppler ultrasound, and equipment controls. This includes colour flow and power Doppler, colour and pulsed wave, scale, gain, filter, angle correction, electronic steering, invert, sample gating, power output, colour amplitude, velocity measurement, spectral changes and all other parameters required to perform a complete diagnostic vascular duplex study.
- Sectional and ultrasonic anatomy
  - Peripheral extremity arteries
  - Peripheral extremity veins
  - Abdominal vessels
  - Extracranial vessels
  - Common normal variants
- Pathology in relation to ultrasound
  - Peripheral extremity arteries: patency, occlusion, stenosis, aneurysmal dilatation
  - Peripheral extremity veins: patency, occlusion, deep venous thrombosis, reflux and incompetence
  - Abdominal vessels: patency, occlusion, aneurysmal dilatation of aorta
  - Extracranial vessels: patency, occlusion, stenosis
  - Appearances and sequelae of common surgical or radiological interventions including angioplasty, stenting, grafts, miller vein cuffs, dissections, and neointimal hyperplasia

### Level 1. Training and practice

Practical training should involve at least two ultrasound lists per week over a period of no less than three months and up to six months, with approximately four to six examinations performed by the trainee under supervision per session.

A minimum of 100 examinations should be undertaken if this is the first practical training module undertaken.

Examinations should encompass the full range of pathological conditions listed below.

A logbook listing the types of examinations undertaken should be kept.

Training should be supervised either by someone who has obtained at least Level 2 competence in vascular ultrasound or by a Level 1 practitioner with at least two years' experience of Level 1 practice. This will usually mean that training is carried out in dedicated vascular duplex sessions supervised by an accredited vascular scientist/technologist, specialist sonographer or radiologist.

Trainees should attend an appropriate theoretical course and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

### **Level 1. Competencies to be acquired**

- To be able to perform continuous wave hand-held Doppler and segmental pressures (ABPI)

#### *Lower extremity peripheral arteries and grafts*

To be able to:

- Perform a complete ultrasound examination of the common femoral to popliteal arteries
- Recognise and assess patency, occlusion, stenosis and aneurysmal dilatation, and measure approximate extent of abnormality
- Diagnose > 50% stenosis (a doubling of peak systolic velocity [PSV] with pulsed Doppler over adjacent segments)
- Recognise common surgical interventions, arteriovenous (AV) fistulas and pseudoaneurysm formation

#### *Peripheral veins*

- Lower extremity deep veins  
To be able to:
  - Perform a complete ultrasound examination of femoral to popliteal deep veins
  - Perform compression and augmentation
  - Recognise acute above-knee deep venous thrombosis
  - Recognise, diagnose and locate reflux
- Lower extremity superficial veins  
To be able to:
  - Identify the saphenofemoral and saphenopopliteal junctions
  - Recognise and locate clinically relevant venous reflux, incompetence and perforators
  - Perform vein mapping and marking
- Abdominal vessels  
To be able to:
  - Recognise and locate patency and occlusion of the abdominal aorta
  - Recognise and size aneurysmal dilatation of the abdominal aorta
- Extracranial vessels  
To be able to:
  - Recognise and locate patency, occlusion, plaque and stenoses in the carotid vessels

### **Level 2. Knowledge base**

- Peripheral arteries and grafts
- Peripheral deep and superficial veins
- Transcranial Doppler ultrasound:
  - Ultrasonic anatomy, common normal variants and principles and practice of the technique
  - Clinical indications and ultrasonic findings in common clinically relevant abnormalities

### **Level 2. Training and practice**

Practical training should include at least one year of experience at Level 1 with continuous ongoing regular ultrasound sessions.

A logbook of all examinations undertaken should be kept.

Supervision of training should be undertaken by someone who has achieved at least Level 1 competence in vascular ultrasound and has had at least two years' experience at that level.

### **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice and refined during a period of clinical practice

To be able to:

- Perform a complete ultrasound scan and identify all abnormalities detailed in Level 1 in the upper and lower extremities, from iliac to infrapopliteal and subclavian to radial and ulnar arteries and veins

### *Extracranial vessels*

To be able to:

- Recognise and diagnose patency, occlusion, stenosis, reverse flow and steal in the carotid and vertebral vessels
- Grade degrees of carotid stenosis and plaque type in accordance with local criteria and standards

### *Abdominal vessels*

To be able to:

- Recognise common normal variants, aneurysmal dilatation, patency, stenosis and occlusion of the major abdominal and iliac vessels, including the mesenteric and renal vessels

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking vascular ultrasound.

They will accept tertiary referrals from Level 1 and 2 practitioners.

They should have the capability to utilise developing technologies and ultrasound techniques, develop research and teaching skills and the performance of specialised examinations including the use of non-invasive physiological studies, contrast agents, intravascular or intra-operative ultrasound and ultrasound-guided invasive procedures.

### **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

A trainee should continue to perform ultrasound scans during the remainder of their training programme, ideally one session weekly and at least 50 examinations per year.

A similar minimum ongoing commitment should be required from a trained practitioner. It is recognised that due to training or clinical circumstances such further ultrasound practice may be intermittent. If a significant period has elapsed after the use of such skills, and a period of re-training is required, it should be agreed and documented with the practitioner, local trainers and assessors.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

The Royal College of Radiologists is grateful to the Vascular Surgical Society of Great Britain and Ireland and the Society of Vascular Technologists who contributed to and approved this section of the document.



# Appendix 5. Breast ultrasound

This curriculum is intended for clinicians who perform diagnostic breast ultrasound and ultrasound-guided breast intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic breast imaging.

## Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasound anatomy
  - Normal anatomy of female and male breast, and patterns of disease spread
  - Changes in ultrasound appearances associated with age, pregnancy and lactation, hormonal status, medication
- Pathology in relation to ultrasound
  - Benign conditions to include: cyst, fibroadenoma, lipoma, fat necrosis, hamartoma, papillary lesions, radial scar/complex sclerosing lesion, gynaecomastia
  - Indeterminate conditions to include: atypical ductal and lobular hyperplasia
  - Malignancy to include: ductal, lobular, inflammatory and other carcinomas
  - Normal and abnormal appearances of axillary lymph nodes
  - Inflammatory breast conditions to include: infection, abscess formation and periductal sepsis
  - Iatrogenic appearances to include: breast implants, early and late postoperative changes, seroma, haematoma, radiotherapy changes, fat necrosis, scarring
- Triple assessment
  - Local/national guidelines
  - Clinical findings associated with normal, benign and malignant changes
  - Mammographic interpretation, correlation and co-localisation with ultrasound findings

## Level 1. Training and practice

Practical training should involve at least one ultrasound session per week over a period of no less than six months and no more than one year.

A minimum of 100 examinations should be undertaken and a record of these kept.

A logbook of 50 cases should be kept which should record details of the indications for the procedure, the interpretation and a final report. These cases should be supported by correlation with clinical examination and other imaging findings, needle biopsy results and surgical histology where appropriate.

The numbers suggested will vary depending how quickly competencies are achieved.

Examinations should include an appropriate range of normal and abnormal cases including palpable and impalpable lesions which should encompass the full range of conditions listed above. They should also include patients presenting to symptomatic clinics, screening assessment clinics and postoperative surgical clinics.

Training should be supervised either by someone who has obtained at least Level 2 competence in breast ultrasound or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should be working in line with National Occupational Standards.<sup>3</sup> The practical experience should ideally be undertaken in conjunction with attendance on a recognised postgraduate course and trainees should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

## Level 1. Competencies to be acquired

- An understanding of the strengths, weaknesses and limitations of breast ultrasound
- An understanding of the indications for and the importance of ultrasound in the triple assessment process
- An awareness of the interdependency and significance of mammographic and ultrasound appearances
- The ability to perform a thorough ultrasound examination of the breast and axilla, to recognise normal anatomy and physiological variation as detailed above and to confidently exclude the presence of a sonographic lesion within the breast
- The recognition of established criteria for lesion characterisation
- Identification and discrimination between clearly normal and suspicious axillary lymph nodes
- The ability to write a detailed report of the ultrasound findings with grading, differential diagnosis, conclusion and recommendation for further management

- The recognition of personal limitations and ability to ask for more expert advice if required

## **Level 2. Knowledge base**

### *Breast imaging*

- Knowledge of a range of *other* imaging studies relevant to breast imaging and their role; eg, breast MR, PET-CT, tomosynthesis, contrast-enhanced mammography, elastography
- An understanding of principles of evaluation of tumour response to treatment

### *Pathology in relation to ultrasound*

- More detailed understanding of breast disease

## **Level 2. Training and practice**

Practical training should involve at least one year of experience at Level 1. Competencies should be further developed during a period of clinical practice, which will involve at least one session per week with at least ten examinations per week for at least three months.

Training for interventional techniques should include observation initially followed by performance of the examination and/or procedure under close supervision. When competence has been acquired, procedures may be undertaken alone but with support close to hand.

A logbook of diagnostic and interventional procedures performed should be kept with pathological correlation.

Supervision of training should be undertaken by someone who has achieved Level 2 competence in breast ultrasound and has had at least two years' experience at that level and who would normally be of consultant status.

The Level 2 practitioner should be competent to accept referrals from Level 1 practitioners.

## **Level 2. Competencies to be acquired**

- Cyst aspiration: initially to perform a minimum of ten guided cyst aspirations of which at least five should be of cysts smaller than 2 cm.
- Aspiration of cysts of less than 1 cm diameter.
- Guided core biopsies of breast: perform a minimum of ten\* guided core biopsies with pathological correlation of core biopsy histology and final pathology (if available).
- Guided fine needle aspiration biopsy (FNAB): perform a minimum of ten\* FNABs of solid lesions, with pathological correlation of cytology result and final pathology (if available).
- Guided core biopsy or FNAB of suspicious axillary lymph nodes, in accordance with local guidelines: perform a minimum of ten\* guided biopsies with correlation of histology/cytology result and final pathology (if available).
- Perform guided abscess aspiration and drainage.
- Perform preoperative guided localisations using skin marking and wire insertion techniques with mammographic corroboration as appropriate.
- Perform guided marker insertion before neo-adjuvant chemotherapy.
- Evaluation of tumour response to neo-adjuvant treatment.
- Evaluation of integrity of breast prostheses.
- Ability to accept referrals from Level 1 practitioners.

\* The numbers suggested may vary according to the practice of individual breast units and the speed with which the trainee achieves competence

## **Level 3. Knowledge base**

- A more detailed understanding of breast imaging and pathology as detailed above including understanding of population screening, risk and prognostic factors as applied to breast cancer.
- Awareness of developments in breast ultrasound including elastography, Doppler and the use of intravascular contrast agents.

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking either all aspects of ultrasound, teaching, research and development or be a specialist breast radiologist and will be an 'expert' in this area.

They will accept tertiary referrals from Level 1 and 2 practitioners and will perform specialised examinations as well as performing advanced ultrasound-guided invasive procedures such as vacuum-assisted breast biopsy.

## **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills. A specialist registrar will need to continue to perform breast ultrasound throughout the remainder of the training programme. Such further ultrasound practice may be intermittent, but no more than three months should elapse without the trainee using their scanning skills.

In medical practice, a clinician scanning at Level 1 should perform at least one ultrasound session per week and a minimum of 150 examinations per year and have regular meetings with radiological colleagues and should have a named radiologist as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

Approved by the British Society of Breast Radiology

## Appendix 5. Breast ultrasound training – competency assessment sheet

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
<b>Competencies/skills to be acquired — Level 1</b>					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
Understands strengths, weaknesses and limitations of breast ultrasound			Aware of relationship between mammographic and ultrasound appearances		
Recognises established criteria for lesion characterisation			Able to write detailed report with grading, differential diagnosis, conclusion and recommendation for further management		
Understands indications for & importance of ultrasound in triple assessment			Can perform thorough ultrasound examination of breast and axilla, recognise normal anatomy and physiological variation and confidently exclude presence of a sonographic lesion		
Identifies and discriminates between clearly normal and suspicious axillary lymph nodes			Knows when to ask for more expert advice		
<b>Competencies/skills to be acquired — Level 2</b>					
To be competent to perform/ diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
Cyst aspiration: initially to perform a minimum of 10 guided cyst aspirations of which at least five should be of cysts smaller than 2 cm			Perform ultrasound-guided breast abscess aspiration		
Aspirate cysts smaller than 1 cm in diameter			Perform ultrasound-guided localisations preoperatively, using both skin marking and wire insertion techniques		
Perform a minimum of 10 FNABs of solid lesions with pathological correlation. If FNA is not standard procedure within the unit then the student should be familiar with the technique and be aware of its uses and limitations			Perform ultrasound-guided marker or coil insertion prior to neo-adjuvant chemotherapy		
Perform a minimum of 10 guided core biopsies with pathological correlation			Accept referrals from Level 1 practitioners		

## Appendix 6. Thoracic ultrasound

The curriculum is intended for clinicians who perform diagnostic and therapeutic ultrasound. At least Level 1 should be obtained by anyone performing thoracic scans unsupervised.

### Focused training and practice

There are frequent situations arising in clinical practice where rapid bedside assessment using focused ultrasound techniques can help with the assessment of, and treatment planning for, patients. In this situation, rapid ultrasound assessment by a competent non-radiological clinician may be more appropriate than waiting for a formal ultrasound list during normal working hours. Some clinicians may wish to focus on only one skill such as ultrasound-guided drainage of pleural effusions. These unitary skills may aid the clinician's practise and greatly improve patient pathways.

Each clinician will have their own requirement for focused training and in order to accommodate their training requirements, a targeted curriculum and syllabus should be created by local trainers, drawing on appropriate elements of the knowledge base and competencies to be acquired from Levels 1–2, depending on the level of practice expected. An example syllabus is included in the focused ultrasound document for example, ultrasound-guided drainage of pleural effusions.<sup>1</sup>

It is essential that ALL ultrasound examinations that may have any influence on patient management are performed by individuals who are competent to provide an accurate examination and assessment and that the images and a formal report are recorded on a RIS/PACS system.

### Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasonic anatomy
  - Right and left hemidiaphragms
  - Heart
  - Liver and spleen
  - Rib and intercostal spaces
- Pathology in relation to ultrasound
  - Pleural effusion
  - Pleural thickening
  - Consolidated lung
  - Paralysed hemidiaphragm
  - Pericardial effusion

### Level 1. Training and practice

Practical training should include:

- At least one session per week over a period of no less than three months, with approximately five scans per session performed by the trainee (under supervision of an experienced practitioner)
- The numbers suggested will vary depending how quickly competencies are achieved.

Examinations should encompass the full range of pathological conditions listed above.

A logbook listing the types of examinations and procedures undertaken should be kept.

Training should be supervised either by someone who has obtained at least Level 2 competence in thoracic ultrasound or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should attend an appropriate theoretical course and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

### Level 1. Competencies to be acquired

- Recognition of normal anatomy of pleura and diaphragm.
- Identification of the heart, liver and spleen.
- Pleural effusion recognition, including the different echogenic patterns.
- Pleural thickening and its differentiation from fluid using colour flow Doppler if appropriate.
- Consolidated lung and its differentiation from effusion.

- Estimation of depth of effusion and its measurement.
- Guided thoracocentesis and drain placement.

## **Level 2. Knowledge base**

### *Sectional and ultrasonic anatomy*

- A full understanding of thoracic and diaphragmatic anatomy

### *Pathology in relation to ultrasound*

- More detailed understanding of pleural disease

## **Level 2. Training and practice**

Practical training should involve at least one year of experience at Level 1 with a minimum of two examinations performed per week.

A further 100 examinations should have been undertaken in order to encompass the full range of conditions and procedures referred to above in addition to those listed below.

A logbook of all examinations undertaken should be kept.

Supervision of training should be by someone who has achieved at least Level 2 competence in thoracic ultrasound, has had at least two years' experience at that level, and who would normally be of consultant status.

A Level 2 practitioner will be able to accept referrals from Level 1 practitioners.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice, and refined during a period of practice.

### *Ultrasound techniques*

- The advanced use of Doppler ultrasound, including spectral, colour and power Doppler
- The use of ultrasound for guiding interventional procedures including lung and supraclavicular nodal biopsy

## **Level 3. Knowledge base**

It is important that physicians correctly interpret CXRs and CT scans used in the decision-making process when managing patients.

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking either all aspects of ultrasound, teaching, research and development or be a specialist thoracic radiologist and will be an 'expert' in this area.

They will accept tertiary referrals from Level 1 and 2 practitioners and will perform specialised examinations as well as performing advanced ultrasound-guided invasive procedures.

## **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills. A chest physician specialist registrar will need to continue to perform ultrasound scans throughout the remainder of the training programme. Such further ultrasound practice may be intermittent, but no more than three months should elapse without the trainee using their scanning skills.

In medical practice, a chest physician scanning at Level 1 should perform at least 20 ultrasound examinations per year and have regular meetings with radiological colleagues and should have a named radiologist as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

## Appendix 6. Thoracic ultrasound training – competency assessment sheet

Core knowledge base – Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
<b>Competencies/skills to be acquired – Level 1</b>					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
Normal anatomy of pleura and diaphragm			Consolidated lung and its differentiation from effusion		
Identification of heart, liver and spleen			Estimation of depth of effusion and its measurement		
Pleural effusion and different echogenic patterns			Know when to refer to a more expert ultrasonologist		
Pleural thickening and its differentiation from fluid					

# Appendix 7. Cranial ultrasound in infants

This syllabus is designed to include the whole of cranial ultrasound imaging in infants, not just the ultrasound skills needed on a neonatal intensive care unit.

It does not include spinal ultrasound imaging.

## Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Issues of parental information and consent.
- Sectional and ultrasonic anatomy of the brain:
  - Sagittal anatomy
  - Coronal anatomy
  - Basic transaxial and near-field scanning
- Pathology in relation to ultrasound:
  - Intracranial haemorrhage
  - Hypoxic ischaemic change (in full-term and premature infants)
  - Ventricular dilatation
  - Common congenital malformations

## Level 1. Training and practice

Practical training should involve at least one session per week over a period of no less than three months, with approximately five scans per session performed by the trainee (under supervision of an experienced practitioner).

The practitioner should be compliant with local procedures for infection control for scanning, especially on the neonatal unit.

By the end of Level 1 training, the trainee should be able to perform and record images from a basic scan demonstrating normal anatomical structures. They should recognise all common pathologies that are seen on a neonatal intensive care unit.

Different trainees will acquire the necessary skills at different rates. The endpoint of the training program should be judged by assessment of competencies, rather than by numbers of scans performed alone. As a guide, a minimum of 100 scans should be undertaken if this is the first practical training module undertaken. For those who are already competent at ultrasound in other body areas, then a reduced number of scans may be needed to achieve competency.

Examinations should encompass all pathological conditions listed below.

A logbook listing the types of examinations undertaken should be kept.

Training should be supervised either by a practitioner who has obtained at least Level 2 competence in cranial ultrasound, or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should attend an appropriate theoretical course (including physics of US and instrumentation) and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

## Level 1. Competencies to be acquired

To be able to:

- Perform a thorough ultrasound examination of the brain in different scan planes
- Recognise normal anatomy and common normal variants
- Recognise varying presentations of pathological processes with gestational and postnatal age
- Measure ventricular size and assess variation from normality
- Assess obstructive hydrocephalus and monitor progression
- Recognise and describe intracranial haemorrhage and assess its extent
- Recognise common congenital malformations and refer for appropriate further investigation
- Recognise when other imaging modalities are more appropriate than ultrasound (eg, MRI or CT)
- Recognise abnormalities which need referral for scanning by a more experienced practitioner and/or further investigation

To be able to use ultrasound in both the routine and urgent assessment of patients presenting with:

- Prematurity
- Fits/apnoea/collapse
- Meningitis
- Hydrocephalus
- Asphyxia (full term and premature)
- Other congenital abnormalities

## **Level 2. Training and practice**

Practical training should involve at least one year of experience at Level 1 with a minimum of one session per week.

A significant number of further examinations should have been undertaken sufficient to encompass the full range of conditions and procedures referred to below.

A logbook of all examinations undertaken should be kept.

Supervision of training should be by a practitioner who has achieved at least Level 2 competence in cranial ultrasound, has had at least two years' experience at that level, and who would normally be of consultant or independent practitioner status.

A Level 2 practitioner will be able to accept referrals from Level 1 practitioners.

## **Level 2. Knowledge base**

### *Sectional and ultrasonic anatomy*

- The basic use of Doppler ultrasound, including spectral, colour and power Doppler
- Further applications of cranial ultrasound
- Use of transaxial ultrasound
- Posterior fossa imaging
- Assessment of cerebral perfusion

### *Pathology in relation to ultrasound*

- Understand of the role of ultrasound in the context of cerebral asphyxia, abnormal head circumference, congenital abnormalities, non-accidental injury and systemic disease and the role of alternative imaging modalities.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice, and refined during a period of practice.

- Vascular studies in asphyxia.
- Role of ultrasound in the assessment of abnormal head circumference.
- Recognition of most identifiable congenital brain malformations.
- Further assessment of antenatally suspected anomalies.
- Identification and location of surface collections.
- Appearances of non-accidental injury and the limitations of ultrasound assessment.
- Intraoperative ultrasound as appropriate.
- Infants with systemic disease (eg, postoperative or paediatric intensive care).

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend a significant proportion of his or her time undertaking cranial ultrasound, teaching, research and development and will be an 'expert' in this area.

They will accept tertiary referrals from Level 1 and 2 practitioners and will perform specialised examinations.

## **Maintenance of skills: all levels**

Having been assessed as competent to practice at Level 1, there will be a need for maintenance of practical skills, by continuing to perform regular ultrasound and update skills. Such further ultrasound practice may be intermittent, but no more than three months should elapse without the trainee using their ultrasound skills, and sufficient examinations should be performed per year to maintain competency.

In independent practice, a medical practitioner scanning at Level 1 should continue to scan regularly should have regular meetings with imaging colleagues and should have a designated ultrasound practitioner of Level 2 experience or above designated as their mentor.

Practitioners should:

- Include ultrasound in their ongoing CPD
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

## Appendix 7. Cranial ultrasound training – competency assessment sheet

Core knowledge base – Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
Infection control					
Competencies/skills to be acquired – Level 1					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
Normal US anatomy of the neonatal brain/variants			Image orientation and documentation		
US assessment of intracranial haemorrhage			US assessment of neonatal asphyxia		
US assessment of ventricular dilatation/large head			US assessment of congenital malformations		
US assessment of the premature infant's brain			Know when to refer to a more expert ultrasonologist		
US assessment of the acutely unwell neonate					

## Appendix 8. Focused emergency ultrasound

This curriculum is intended for clinicians who perform diagnostic ultrasound and ultrasound-guided intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

### Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasonic anatomy
  - Kidneys
  - Liver
  - Spleen
  - Retroperitoneal structures (aorta, inferior vena cava [IVC])
  - Vessels: internal jugular veins, carotid arteries, femoral veins and arteries
- Pathology in relation to ultrasound
  - Kidneys: trauma/free fluid
  - Liver and spleen: trauma/free fluid
  - Retroperitoneal: presence or absence of abdominal aortic aneurysm (AAA)
  - Vessels: vascular access
  - Cardiac scan: trauma/pericardial tamponade, pericardial effusions, asystole

### Level 1. Training and practice

Practical training should involve regular emergency department or radiology department ultrasound, with approximately five examinations performed by the trainee (under supervision) per week.

Approximately 50 examinations should be undertaken if this is the first practical training module. A clinician working in emergency medicine will need to devote sufficient time to gain Level 1 competence. This may be as much as one session per week but is unlikely to be in dedicated blocks of time. However, different trainees will acquire the necessary skills at different rates, and the key endpoint of the training programme should be judged by an assessment of competencies.

Examinations should concentrate on the core clinical indications of trauma, aortic aneurysm and vascular access where there are benefits of an early focused ultrasound scan in the emergency department or acute assessment area.

A logbook listing the types of examinations undertaken should be kept.

An additional pictorial record containing an illustrated description of ten cases in which the trainee has been personally involved may be collected and is a useful confirmation of experience when moving between departments.

Training should be supervised either by someone who has obtained at least Level 2 competence in focused emergency ultrasound, or by a Level 1 practitioner with at least one year's experience of Level 1 practice.

Trainees should attend an appropriate theoretical course and should be familiar with the published literature on focused emergency ultrasound.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

### Level 1. Competencies to be acquired

To be able to:

- Recognise normal anatomy
- Use focused ultrasound to assist in bedside emergency department decisions
- Recognise the limitations of a scan and be able to explain these limitations to patients/carers
- Recognise patients requiring formal specialist sonographic assessment
- Incorporate ultrasound findings with the rest of the clinical assessment

To be able to use ultrasound in the context assessment of:

- Focused assessment by sonography for trauma (FAST)
- AAA screening/detection in symptomatic patients
- Peri-arrest scenario for pulseless electrical activity (PEA)/tamponade/effusion

- Vascular access
- Pleural and pericardial fluid

## **Level 2. Knowledge base**

- Sectional and ultrasonic anatomy
  - Kidneys
  - Liver
  - Proximal leg veins
  - Heart
  - Musculoskeletal system
  - Pelvic structures (uterus, ovaries, bowel)
- Pathology in relation to ultrasound
  - Kidneys: cysts, calyceal dilatation, renal calculi, trauma (free fluid)
  - Liver: cysts, trauma (free fluid)
  - Vessels: proximal deep venous thrombosis
  - Cardiac: reduced ventricular function, tamponade, effusion
  - Retroperitoneum: aortic aneurysm, fluid collection
  - Musculoskeletal: fractures, soft tissue collections, foreign body detection

## **Level 2. Training and practice**

Practical training should involve at least one year of experience at Level 1 with an average of three to five scans/week.

A further 150–200 examinations should have been undertaken in order to encompass most of the conditions and procedures listed. A practitioner may develop competencies in some but not all of these areas.

A logbook of all examinations undertaken should be kept.

A further pictorial logbook should also be kept detailing ten cases examined by the trainee which may be useful as a record when moving between departments.

Supervision of training should be undertaken by someone who has achieved at least Level 2 competence in focused emergency ultrasound and who has had at least two years' experience at this level.

A Level 2 practitioner will be able to accept referrals from a Level 1 practitioner.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice, and refined during a period of clinical practice.

- Recognise and correctly utilise ultrasound within the emergency department and be able to perform at least three of the following:
  - Ultrasound-guided invasive procedures (chest drain insertion, suprapubic aspiration/ catheterisation, fluid collection drainage)
  - An initial assessment for patients with loin pain/haematuria
  - A focused assessment of a patient presenting with hypotension
  - Proximal DVT assessment
  - Detection of foreign bodies and fluid collections within soft tissues
  - Early assessment of symptomatic women in the first trimester of pregnancy post-trauma including assessment of foetal cardiac activity
  - Emergency department obstetric presentations

## **Level 3. Training and practice**

A Level 3 practitioner is likely to spend a significant proportion of their time undertaking emergency ultrasound, teaching, research and development and will be an 'expert' in this area.

They will accept referrals from Level 1 and 2 practitioners and will perform the spectrum of examinations indicated above. This will include developing the role for innovative ultrasound practice within emergency care.

## **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

An emergency trainee will need to continue to perform ultrasound scans throughout the remainder of the training programme and into their consultant appointment. Such further ultrasound practice may be intermittent, but no more than three months should elapse without the trainee using his scanning skills, and at least 50 scans should be performed per year.

All practitioners should have regular meetings within the department to ensure appropriate focused emergency ultrasound use. The department lead for ultrasound practice will have regular contact with radiological colleagues and should have a named radiologist as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature

Approved by the Faculty of Accident and Emergency Medicine

## Appendix 8. Focused emergency ultrasound training – competency assessment sheet

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
<b>Competencies/skills to be acquired — Level 1</b>					
To be competent to perform/diagnose/recognise the following:					
	Trainer signature	Date		Trainer signature	Date
Normal anatomy and variants of kidneys, liver, spleen and aorta			Ultrasound in the assessment of patients presenting with:		
Differences between arteries and veins			<ul style="list-style-type: none"> <li>▪ Trauma-FAST</li> </ul>		
Focused ultrasound to assist in bedside emergency			<ul style="list-style-type: none"> <li>▪ AAA screening/detection in symptomatic patients</li> </ul>		
Department decisions			<ul style="list-style-type: none"> <li>▪ Peri-arrest scenario for PEA/tamponade/effusion</li> </ul>		
The limitations of a scan and be able to explain these limitations to patients/carers			<ul style="list-style-type: none"> <li>▪ Vascular access: internal jugular and femoral vein</li> </ul>		
Patients requiring formal specialist sonographic assessment			Pleural and pericardial fluid		
Ultrasound findings with the rest of the clinical assessment			<ul style="list-style-type: none"> <li>▪ Free intraperitoneal fluid – use of four-quadrant scan</li> </ul>		
Core knowledge base — Level 2		Trainee		Trainer	
	Trainer signature	Date		Trainer signature	Date
Sectional and ultrasonic anatomy			Pathology in relation to ultrasound		
<ul style="list-style-type: none"> <li>▪ Kidneys</li> </ul>			<ul style="list-style-type: none"> <li>▪ Kidneys: cysts, calyceal dilatation, renal calculi, trauma (free fluid)</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Liver</li> </ul>			<ul style="list-style-type: none"> <li>▪ Liver: cysts, trauma (free fluid)</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Proximal leg veins</li> </ul>			<ul style="list-style-type: none"> <li>▪ Vessels: proximal deep venous thrombosis</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Heart</li> </ul>			<ul style="list-style-type: none"> <li>▪ Cardiac: reduced ventricular function, tamponade, effusion</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Musculoskeletal system</li> </ul>			<ul style="list-style-type: none"> <li>▪ Retroperitoneum: aortic aneurysm, fluid collection</li> </ul>		
<ul style="list-style-type: none"> <li>▪ Pelvic structures (uterus, ovaries, bowel)</li> </ul>			<ul style="list-style-type: none"> <li>▪ Musculoskeletal: fractures, soft tissue collections, foreign body detection</li> </ul>		
<b>Competencies/skills to be acquired — Level 2</b> Recognise the role of and correctly utilise ultrasound within the emergency department and be able to perform at least three of the following:					
	Trainer signature	Date		Trainer signature	Date
Ultrasound-guided invasive procedures (chest drain insertion, suprapubic aspiration/catheterisation, fluid collection drainage)			Detection of foreign bodies and fluid collections within soft tissues		
An initial assessment for patients with loin pain/haematuria			Early assessment of symptomatic women in the first trimester of pregnancy after trauma including the assessment of foetal cardiac activity		
A focused assessment of a patient presenting with hypotension			Emergency department obstetric presentations		
Proximal DVT assessment					

## Appendix 9. Ultrasound training for critical/intensive care

This curriculum is intended for clinicians who perform diagnostic ultrasound and ultrasound-guided intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

### Level 1. Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasonic anatomy
  - Thorax
  - Abdomen, retroperitoneal and peritoneal spaces
  - Abdominal aorta
  - Urinary tract
  - Peripheral vascular anatomy
  - Peripheral neurological anatomy
- Pathology in relation to ultrasound
  - Thorax: pleural disease, patterns and causes of pulmonary consolidation, pericardial disease.
  - Abdomen: fluid collections, gastrointestinal ileus, acute urinary tract pathology, aortic aneurysms
  - Vascular: venous disease, eg, thrombosis
- Basic knowledge of Doppler ultrasound principles

### Level 1. Training and practice

Practical training should involve carrying out regular ultrasound examinations in the critical care unit or radiology department, with approximately 5–10 examinations performed by the trainee (under supervision) per week.

A minimum number of examinations in each specific area of interest should be undertaken if this is the first practical training module undertaken; for example, 25 pleural effusions, 25 vascular line insertions, 25 abnormal abdominal scans. However, different trainees will acquire the necessary skills at different rates and the endpoint of the training programme should be judged by an assessment of competencies.

Examinations should encompass the full range of pathological conditions and practical procedures listed below.

A logbook listing the types of examinations undertaken should be kept.

An additional logbook containing an illustrated description of ten cases in which the trainee has been personally involved may be a useful adjunct particularly if moving between different units.

Training should be supervised either by someone who has obtained at least Level 2 competence in critical care ultrasound themselves, or by a Level 1 practitioner with at least two years' experience of Level 1 practice.

Trainees should attend an appropriate theoretical course and should be familiar with the published literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

### Level 1. Competencies to be acquired

#### *General*

- Recognise normal anatomy
- Use ultrasound to assist in critical care decisions and interventional procedures
- Recognise limitations of technique and own knowledge base
- Recognise patients requiring formal specialist sonographic assessment

#### *Thorax*

- Pleural disease
  - Identify and assess pleural effusions: transudate, exudates, volume estimation
  - Use ultrasound to guide pleural aspiration/drainage
- Pulmonary
  - Identify pulmonary consolidation
  - Distinguish abnormal lung from pleural diseases
- Pericardial
  - identify significant pericardial effusions prior to formal echocardiographic assessment

### *Abdomen*

- Fluid collections
  - Identify and assess abdominal and pelvic fluid collections
  - Differentiate subphrenic from pleural fluid
- Distension
  - Identify cause of abdominal distension, eg, fluid, small bowel ileus, bladder distension, masses

### *Urinary tract*

- Identify cause of acute renal impairment, specifically hydronephrosis
- Identify significant bladder distension

### *Vascular*

- Identify and measure abdominal aortic aneurysm
- Recognise abdominal features secondary to cardiac failure

### *Peripheral vascular*

- Differentiate between arteries and veins
- Assess veins for vascular access
  - Identify suitable neck veins and their patency
  - Use ultrasound to guide line insertions

### *Peripheral neurology*

- Identify paths of appropriate peripheral nerves, eg, brachial plexus
- Use ultrasound to guide regional anaesthesia

## **Level 2. Knowledge base**

- Physics and technology
  - In-depth knowledge and understanding of physics of ultrasound
  - In-depth knowledge and understanding of the technology of ultrasound equipment
- Ultrasound techniques
  - Further applications of thoracic, abdominal, vascular and cardiac ultrasound
- Sectional and ultrasonic anatomy
  - More detailed recognition and understanding of normal variants with ultrasound
- Pathology related to ultrasound

## **Level 2. Training and practice**

Practical training should include at least one year of experience at Level 1 with a minimum of one session or equivalent per week.

A further 300 examinations should have been undertaken in order to encompass the full range of conditions and procedures.

A logbook listing all examinations undertaken should be kept.

A further illustrated logbook should be kept detailing 20 cases examined by the trainee which may be useful to document further progression of training.

Supervision of training should be undertaken by someone who has achieved at least Level 2 competence in critical care ultrasound, has had at least two years' experience at that level and would normally be of consultant status.

A Level 2 practitioner will be able to accept referral from Level 1 practitioners.

## **Level 2. Competencies to be acquired**

Competencies will have been gained during training for Level 1 practice, and refined during a period of clinical practice.

- Develop further ultrasound skills such as echocardiography, transoesophageal echocardiography and perioperative monitoring.

### **Level 3. Training and practice**

A Level 3 practitioner is likely to spend the majority of their time undertaking ultrasound examinations, teaching, research and development, and will be regarded as an 'expert' in this area.

They will accept referrals from Level 1 and 2 practitioners and will be actively involved in developing innovative ultrasound applications within critical care.

They will have spent a continuous period of specialist training in critical care ultrasound.

They will be able to mentor and supervise Level 1 and 2 practitioners.

They will be aware of and pursue developments in critical care ultrasound including Doppler and the use of intravascular contrast agents.

#### **Maintenance of skills: all levels**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

A trainee should continue to practise ultrasound throughout their training with no more than one month elapsing without the trainee using their ultrasound scanning skills.

In medical practice, a critical care physician performing ultrasound at Level 1 should continue to perform at least 100 examinations per year, should have regular meetings with radiological colleagues and should have a named radiologist or experienced sonographic practitioner as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary team meetings
- Keep up to date with relevant literature.

## Appendix 9. Critical care ultrasound training – competency assessment sheet

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
<b>Competencies/skills to be acquired — Level 1</b>					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Thorax</b>			<ul style="list-style-type: none"> <li>Understand likely sites of fluid collection</li> </ul>		
<ul style="list-style-type: none"> <li>Normal aerated lung</li> </ul>			<ul style="list-style-type: none"> <li>Differentiate abdominal from pleural fluid</li> </ul>		
<ul style="list-style-type: none"> <li>Pleural effusion</li> </ul>			<ul style="list-style-type: none"> <li>Normal bowel pattern</li> </ul>		
<ul style="list-style-type: none"> <li>Differentiate transudate from exudate</li> </ul>			<ul style="list-style-type: none"> <li>Patterns of ileus</li> </ul>		
<ul style="list-style-type: none"> <li>Estimate volume of pleural fluid</li> </ul>			<ul style="list-style-type: none"> <li>Bladder distension</li> </ul>		
<ul style="list-style-type: none"> <li>Pulmonary consolidation</li> </ul>			<ul style="list-style-type: none"> <li>Normal kidney</li> </ul>		
<ul style="list-style-type: none"> <li>Pericardial effusion</li> </ul>			<ul style="list-style-type: none"> <li>Hydronephrosis</li> </ul>		
<ul style="list-style-type: none"> <li>Use ultrasound to guide pleural aspiration/drainage</li> </ul>			<ul style="list-style-type: none"> <li>Normal aorta</li> </ul>		
<ul style="list-style-type: none"> <li>Identify pulmonary consolidation</li> </ul>			<ul style="list-style-type: none"> <li>Identify and measure aortic aneurysm</li> </ul>		
<b>Vascular</b>			<ul style="list-style-type: none"> <li>Recognise abdominal signs of cardiac failure</li> </ul>		
<ul style="list-style-type: none"> <li>Basic knowledge of Doppler ultrasound principles</li> </ul>			<b>Peripheral neurology</b>		
<ul style="list-style-type: none"> <li>Differentiate between arteries and veins</li> </ul>			<ul style="list-style-type: none"> <li>Recognise paths of appropriate nerves</li> </ul>		
<ul style="list-style-type: none"> <li>Venous patency</li> </ul>			<ul style="list-style-type: none"> <li>Use ultrasound to guide regional anaesthesia</li> </ul>		
<ul style="list-style-type: none"> <li>Assess veins for cannulation</li> </ul>			<b>General</b>		
<ul style="list-style-type: none"> <li>Use ultrasound to guide venous line insertion</li> </ul>			<ul style="list-style-type: none"> <li>Know when to refer to a more expert ultrasonologist</li> </ul>		
<b>Abdomen</b>			<ul style="list-style-type: none"> <li>Understand likely sites of fluid collection</li> </ul>		
<ul style="list-style-type: none"> <li>Fluid collections</li> </ul>					
<b>Core knowledge base — Level 2</b>		<b>Trainee</b>		<b>Trainer</b>	
	Trainer signature	Date		Trainer signature	Date
<b>Physics and technology</b>			<b>Sectional and ultrasonic anatomy</b>		
<ul style="list-style-type: none"> <li>In-depth knowledge and understanding of physics of ultrasound</li> </ul>			<ul style="list-style-type: none"> <li>More detailed recognition and understanding of normal variants with ultrasound</li> </ul>		
<ul style="list-style-type: none"> <li>In-depth knowledge and understanding of the technology of ultrasound equipment</li> </ul>			<b>Pathology related to ultrasound</b>		
<b>Ultrasound techniques</b>					
<ul style="list-style-type: none"> <li>Further applications of thoracic, abdominal, vascular and cardiac ultrasound</li> </ul>					
<b>Competencies/skills to be acquired — Level 2</b>					
To be competent to perform/ diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
Development of further ultrasound skills such as echocardiography, transoesophageal echocardiography and perioperative monitoring					

# Appendix 10. Musculoskeletal ultrasound

This curriculum is intended for clinicians who perform diagnostic ultrasound and ultrasound-guided intervention. At least Level 1 should be obtained by anyone performing unsupervised diagnostic imaging.

Musculoskeletal ultrasound comprises a wide range of examinations and pathologies. It is unlikely that any practitioner will cover the full range. A modular approach that meets the needs of an individual's clinical practice is therefore likely to be appropriate and 'levels' of competence are not stipulated in this section.

## Knowledge base

- Physics and technology, ultrasound techniques and administration (see Appendix 11)
- Sectional and ultrasonic anatomy
  - Shoulder
  - Elbow
  - Wrist/hand
  - Groin
  - Hip
  - Thigh
  - Knee
  - Lower leg
  - Ankle/foot
  - Other (eg, infant spine, brachial plexus)
- Pathology in relation to ultrasound
  - Shoulder: rotator cuff tendinosis, partial and complete thickness tears and calcification; joint and bursal effusion; long head of biceps tendinosis, rupture and subluxation; paralabral cyst
  - Elbow: tendinosis of common flexor/extensor origins; tendinosis and rupture of triceps and distal biceps tendons; effusion; loose bodies; ulnar nerve entrapment; bursae
  - Wrist/hand: tendon tears and tenosynovitis; carpal tunnel syndrome; space-occupying lesions; pulley injuries; arthropathy
  - Groin: hernias; tendinosis and tendon tears
  - Hip: developmental dysplasia of the hip (DDH); effusion (children, adults, prosthetic joints); bursae; snapping hip; tendinosis and tendon tears
  - Thigh: muscle contusion and tears; tendinosis and tendon tears
  - Knee: effusion; bursae (including ruptured baker's cyst); meniscal cysts; collateral ligament sprains; patellar tendon tendinosis and rupture
  - Lower leg: muscle contusion and tears, muscle herniae
  - Ankle/foot: tendinosis and tears of Achilles, posterior tibial and peroneal tendons; tenosynovitis; ankle effusion and loose bodies; plantar fasciitis; Morton's neuroma; other space occupying lesions; arthropathy
  - Soft tissue masses
  - Brachial plexus: injuries and tumours
  - Spine: dysraphism and tethered cord in neonates

## Training and practice

Practical training should involve at least one ultrasound list per week for 3–6 months with approximately ten examinations performed by the trainee under supervision per session.

A minimum of 250 examinations should be undertaken. However, trainees acquire the necessary skills at different rates and some examinations (eg, shoulder ultrasound) take considerably longer than others for trainees to acquire proficiency. The end point of the training programme should be judged by an assessment of competencies.

A logbook listing the types and numbers of examinations undertaken should be kept.

Training should be supervised by an experienced musculoskeletal sonologist.

Trainees should attend an appropriate theoretical course and should read appropriate textbooks and literature.

During the course of training, the competency assessment sheet should be completed as this will determine in which area or areas the trainee can practise independently.

## Competencies to be acquired

### *Other diagnostic techniques*

- Be aware of alternative diagnostic methods including clinical examination and imaging techniques
- Recognise comparative accuracy of alternative techniques
- Recognise when to proceed to other imaging examinations following ultrasound examination

### *Shoulder*

To be able to:

- Perform a thorough ultrasound examination of the shoulder in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Recognise and be aware of difficulties in distinguishing accurately between tendinosis/ partial thickness/ complete thickness tears of the rotator cuff
- Recognise rotator cuff calcification
- Recognise tendinosis, rupture and subluxation of the long head of biceps tendon
- Recognise effusions of the shoulder joint and subdeltoid bursa
- Recognise paralabral cyst
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Elbow*

To be able to:

- Perform a thorough ultrasound examination of the elbow in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Recognise tendinosis of the common flexor/extensor origins
- Recognise tendinosis/partial/complete rupture of the triceps and distal biceps tendons
- Recognise joint effusions and loose bodies
- Recognise ulnar nerve entrapment
- Recognise olecranon bursitis
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Wrist/hand*

To be able to:

- Perform a thorough ultrasound examination in different planes
- Recognise normal ultrasonic anatomy and common normal variants including accessory muscles
- Recognise de Quervain's tenosynovitis
- Recognise effusions in other tendons sheaths
- Recognise tendonitis, partial and complete tendon tears and identify retracted tendon
- Recognise features of carpal tunnel syndrome and entrapment in Guyon's canal
- Recognise ganglia and distinguish them from solid space occupying lesions
- Recognise inflammatory arthropathy
- Recognise pulley injuries
- Recognise foreign bodies (FB) and FB reactions
- Recognise ligament injuries
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Groin*

To be able to:

- Perform a thorough ultrasound examination in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Recognise inguinal and femoral hernias
- Recognise strains of the rectus abdominis and adductor muscles
- Recognise abnormalities which need referral for scanning by a more experienced ultrasonologist and/or further investigation

## *Hip*

To be able to:

- Perform a thorough ultrasound examination in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Assess the neonatal hip and recognise DDH
- Recognise joint effusions in children, adults and prosthetic joints
- Recognise iliopsoas bursa and distinguish from solid masses
- Recognise causes of snapping hip such as iliopsoas, tensor fascia lata and rectus femoris tendons
- Recognise tendinosis and tendon tears eg, of hamstrings, glutei
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

## *Thigh*

To be able to:

- Perform a thorough ultrasound examination in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Recognise contusions and tears of the quadriceps and hamstring muscles
- Recognise tendinosis and partial and complete tears of the hamstring and quadriceps tendons
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

## *Knee*

To be able to:

- Perform a thorough ultrasound examination in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Recognise joint effusion
- Recognise ganglia and bursae, including ruptured Baker's cyst
- Recognise meniscal cysts and associated meniscal tears
- Recognise sprains of collateral ligaments
- Recognise tendinosis and tears of patellar tendon
- Recognise quadriceps tendon
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

## *Lower leg*

To be able to:

- Perform a thorough ultrasound examination of the lower leg in different planes
- Recognise normal ultrasonic anatomy and common normal variants
- Recognise muscle contusions and tears
- Recognise muscle herniae

## *Ankle/foot*

To be able to:

- Perform a thorough ultrasound examination in different planes
- Recognise normal ultrasonic anatomy and common normal variants including accessory muscles
- Recognise tendinosis and tears of Achilles, posterior tibial and peroneal tendons
- Recognise joint effusions and loose bodies
- Recognise plantar fasciitis
- Recognise Morton's neuroma
- Recognise ganglia and distinguish them from solid space occupying lesions
- Recognise arthropathy
- Recognise ankle ligament injuries
- Recognise foreign bodies and foreign body reactions
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Soft tissue masses*

To be able to:

- recognise solid, cystic and indeterminate masses
- recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Brachial plexus*

To be able to:

- Recognise brachial plexus injuries and tumours
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Spine*

To be able to:

- Recognise spinal dysraphism and tethered cord in neonates
- Recognise abnormalities which need referral to a more experienced ultrasonologist and/or for further investigation

### *Technique*

To be able to:

- Perform Doppler ultrasound examinations relevant to the MSK system, including spectral, colour and power Doppler

### *Interventional*

To be able to:

- Perform ultrasound-guided invasive procedures including joint and cyst aspiration and injection; abscess drainage; biopsy of soft tissue masses and synovium; therapeutic injections

### **Maintenance of skills**

Having been assessed as competent to practise, there will be a need for CPD and maintenance of practical skills.

A specialist registrar will need to continue to perform ultrasound scans throughout the remainder of their training programme.

Such further ultrasound practice may be intermittent but no more than three months should elapse without the trainee using their ultrasound skills and at least 100 examinations should be performed per year.

A medical practitioner should perform at least 200 examinations per year on a regular basis, should have regular meetings with radiological colleagues and should have a named radiologist as an 'ultrasound mentor'.

Practitioners should:

- Include ultrasound in their ongoing CPD which should be included in annual appraisal and revalidation
- Audit their practice
- Participate in multidisciplinary meetings
- Keep up to date with relevant literature.

Approved by the British Society of Skeletal Radiology

# Appendix 10. Musculoskeletal ultrasound training – competency assessment sheet

Core knowledge base — Level 1		Trainee:		Trainer:	
	Trainer signature	Date		Trainer signature	Date
Physics and technology			Administration		
Practical instrumentation/use of ultrasound controls			Sectional and ultrasonic anatomy		
Ultrasound techniques			Pathology in relation to ultrasound		
<b>Competencies/skills to be acquired — Level 1</b>					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Shoulder</b>			<b>Wrist/hand cont</b>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Normal anatomy and variants</li> <li>▪ Rotator cuff tendinosis, tears and calcification</li> <li>▪ Tendinosis, rupture and subluxation of long head of biceps tendon</li> <li>▪ Effusions of shoulder joint and subdeltoid bursa</li> <li>▪ Paralabral cyst</li> <li>▪ Aspiration of joint or bursal effusion</li> <li>▪ Aspiration of focus of calcification</li> <li>▪ Therapeutic/diagnostic injection in bursa and joint</li> </ul>			<ul style="list-style-type: none"> <li>▪ Pulley injuries</li> <li>▪ Ligament injuries</li> <li>▪ Foreign bodies and reactive changes</li> <li>▪ Ligament injuries</li> <li>▪ Aspiration and therapeutic injections of joints and tendon sheaths</li> </ul>		
<b>Elbow</b>			<b>Groin</b>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Normal anatomy and variants</li> <li>▪ Tendinosis or rupture of common flexor/extensor origins or biceps/triceps tendons</li> <li>▪ Joint effusion and loose bodies</li> <li>▪ Ulnar nerve entrapment</li> <li>▪ Olecranon bursitis</li> <li>▪ Aspiration of joint and bursal effusion</li> <li>▪ Therapeutic/diagnostic injection of joint</li> </ul>			<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Normal anatomy and variants</li> <li>▪ Inguinal and femoral hernias</li> <li>▪ Strains of rectus abdominis and adductor muscles</li> <li>▪ Therapeutic injections</li> </ul>		
<b>Wrist/hand</b>			<b>Hip</b>		
<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Normal anatomy and variants</li> <li>▪ de Quervain's tenosynovitis</li> <li>▪ Effusions in tendon sheaths</li> <li>▪ Tendinosis and tendon ruptures</li> <li>▪ Carpal tunnel syndrome and ulnar nerve entrapment</li> <li>▪ Ganglia and distinguish them from solid space-occupying lesions</li> <li>▪ Inflammatory arthropathy</li> </ul>			<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Normal anatomy and variants at different ages</li> <li>▪ DDH</li> <li>▪ Joint effusions</li> <li>▪ Iliopsoas bursa</li> <li>▪ Causes of snapping hip</li> <li>▪ Tendinosis and tears of glutei and hamstrings</li> <li>▪ Joint aspiration and therapeutic injections</li> <li>▪ Bursal aspiration and therapeutic injection</li> </ul>		
			<b>Thigh</b>		
			<ul style="list-style-type: none"> <li>▪ Ultrasound examination in different planes</li> <li>▪ Normal anatomy and variants</li> <li>▪ Contusions and tears of the quadriceps and hamstring</li> <li>▪ Muscles and tendons</li> <li>▪ Aspiration of thigh haematoma</li> </ul>		

Competencies/skills to be acquired — Level 1					
To be competent to perform/diagnose the following:					
	Trainer signature	Date		Trainer signature	Date
<b>Knee</b>			<b>Soft tissue masses</b>		
▪ Ultrasound examination in different planes			▪ Solid, cystic and indeterminate masses		
▪ Normal anatomy and variants			▪ Need for appropriate additional imaging examinations		
▪ Joint effusion			▪ Appropriate aspiration and biopsy		
▪ Baker's cyst and ruptured Baker's cyst			▪ Brachial plexus		
▪ Ganglia/other bursae/meniscal cyst			▪ Normal anatomy and variants		
▪ Sprains of collateral ligaments			▪ Injuries and tumours		
▪ Tendinosis and tears of patellar tendon			<b>Spine</b>		
<b>Lower leg</b>			▪ Normal neonatal anatomy		
▪ Ultrasound examination in different planes			▪ Spinal dysraphism and tethered cord		
▪ Normal anatomy and variants			▪ Technique		
▪ Muscle contusions and tears			▪ Doppler ultrasound examinations relevant to MSK system		
▪ Muscle herniae			▪ Interventional		
<b>Ankle/foot</b>			▪ Joint and cyst aspiration and injection		
▪ Ultrasound examination in different planes			▪ Abscess aspiration and drainage		
▪ Normal anatomy and variants			▪ Biopsy of soft tissue masses		
▪ Tendinosis and tears of tendon Achilles, posterior tibial and peroneal tendons			▪ Synovial biopsy		
▪ Joint effusions and loose bodies			▪ Therapeutic injections		
▪ Plantar fasciitis			<b>General</b>		
▪ Morton's neuroma			▪ Know when to refer to a more expert ultrasonologist		
▪ Ganglia					
▪ Arthropathy					
▪ Ankle ligament injuries					
▪ Foreign bodies					
▪ Joint aspiration and therapeutic injection					
▪ Injection of plantar fasciitis and Morton's neuroma					

# Appendix 11. Recommended theory syllabus

This basic theoretical training is a prerequisite to any practical training in ultrasound.

## Physics and instrumentation

- The basic components of an ultrasound system
- Types of transducer and the production of ultrasound, with an emphasis on operator-controlled variables
- Use of ultrasound controls
- An understanding of the frequencies used in medical ultrasound and the effect on image quality and penetration
- The interaction of ultrasound with tissue including biological effects
- The safety of ultrasound and of ultrasound contrast agents
- The basic principles of real-time and Doppler ultrasound including colour flow and power Doppler
- The recognition and explanation of common artefacts
- Image recording systems

## Ultrasound techniques

- Patient information and preparation
- Indications for examinations
- Relevance of ultrasound to other imaging modalities
- The influence of ultrasound results on the need for other imaging
- Scanning techniques including the use of spectral Doppler and colour Doppler

## Administration

- Image and report recording
- Image and report storing and filing (PACS/RIS)
- Image reporting
- Medico-legal aspects – outlining the responsibility to practise within specific levels of competence and the requirements for training
- Consent
- The value and role of departmental protocols
- The resource implications of ultrasound use

Citation details:

The Royal College of Radiologists. *Ultrasound training recommendations for medical and surgical specialties, Second edition*. London: The Royal College of Radiologists, 2012.

Ref No. BFCR(12)17 © The Royal College of Radiologists, December 2012

For permission to reproduce any of the content contained herein, please email: [permissions@rcr.ac.uk](mailto:permissions@rcr.ac.uk)

This material has been produced by The Royal College of Radiologists (RCR) for use internally within the specialties of clinical oncology and clinical radiology in the United Kingdom. It is provided for use by appropriately qualified professionals, and the making of any decision regarding the applicability and suitability of the material in any particular circumstance is subject to the user's professional judgement.

While every reasonable care has been taken to ensure the accuracy of the material, RCR cannot accept any responsibility for any action taken, or not taken, on the basis of it. As publisher, RCR shall not be liable to any person for any loss or damage, which may arise from the use of any of the material. The RCR does not exclude or limit liability for death or personal injury to the extent only that the same arises as a result of the negligence of RCR, its employees, Officers, members and Fellows, or any other person contributing to the formulation of the material.